For Research Use Only

NARS Monoclonal antibody

Catalog Number: 67711-1-Ig



Purification Method:

Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:2000-1:8000

IF 1:400-1:1600

Basic Information

Applications

Catalog Number: GenBank Accession Number:

Protein A purification 67711-1-lg BC001687 GeneID (NCBI): CloneNo.: 150ul, Concentration: 1000 µg/ml by 4677 4G5G12

Nanodrop and 500 µg/ml by Bradford Full Name:

method using BSA as the standard; asparaginyl-tRNA synthetase

Calculated MW: Mouse 63 kDa Isotype: Observed MW: lgG2a 63 kDa

Immunogen Catalog Number:

Tested Applications:

AG6656

IF, IHC, WB, ELISA WB: LNCaP cells, HeLa cells, HepG2 cells, Jurkat cells, Species Specificity: HSC-T6 cells, NIH/3T3 cells, RAW 264.7 cells

Human, mouse, rat IHC: human urothelial carcinoma tissue,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

IF: HeLa cells,

Background Information

Asparagine-tRNA ligase (NARS) belongs to the class-II aminoacyl-tRNA synthetase family. Catalyzes the attachment of asparagine to tRNA(Asn) in a two-step reaction: asparagine is first activated by ATP to form Asn-AMP and then transferred to the acceptor end of tRNA(Asn) (PubMed: 9421509). In addition to its essential role in protein synthesis, acts as a signaling molecule that induced migration of CCR3-expressing cells. The NARS levels were higher in tumor tissues compared with adjacent normal tissues, and positively associated with LN metastasis. NARS may serve as a potential biomarker for lung adenocarcinoma (ADC) diagnosis/prognosis (PMID: 27161446).

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67711-1-1g (NARS antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed Hela cells using NARS antibody (67711-1-1g, Clone: 4G5G12) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffinembedded human urothelial carcinoma tissue slide using 67711-1-1g (NARS antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).