For Research Use Only

Alix Monoclonal antibody

Catalog Number:67715-1-lg Featured Product

12 Publications



Basic Information

Catalog Number: GenBank Accession Number: BC020066

67715-1-lg Protein A purification GeneID (NCBI): Size: CloneNo.:

150ul , Concentration: 1000 ug/ml by 10015 1H9D9

Nanodrop and 373 ug/ml by Bradford UNIPROT ID: Recommended Dilutions: method using BSA as the standard; Q8WUM4 WB 1:2000-1:20000 Source: IHC 1:4000-1:16000 Full Name:

Mouse IF/ICC 1:200-1:800 programmed cell death 6 interacting

Isotype: protein

lgG1 Calculated MW: 868 aa, 96 kDa Immunogen Catalog Number:

AG30437 Observed MW: 100 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA

Cited Applications:

Species Specificity: human, mouse, rat Cited Species:

human, mouse, rat, monkey

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HSC-T6 cells, A431 cells, HEK-293 cells, HeLa cells, HepG2 cells, Jurkat cells, K-562 cells, PC-12 cells,

Purification Method:

NIH/3T3 cells, 4T1 cells

IHC: human stomach cancer tissue,

IF/ICC: HeLa cells,

Background Information

ALG-2-interacting protein 1 (ALIX), also known as AIP1 or Hp95, is encoded by PDCD6IP gene and is involved in cell death through mechanisms involving its binding partner ALG-2 (apoptosis-linked gene-2). ALG-2 is a 22-kDa protein containing five serially repetitive EF-hand structures and is defined as a regulator of calcium-induced apoptosis following endoplasmic reticulum (ER) stress. ALIX interacts with ALG-2 through its C-terminal proline-rich region and participates in formation of multivesicular bodies. Recent finding suggest that ALIX is a critical component of caspase 9 activation and apoptosis triggered by calcium.

Notable Publications

Author	Pubmed ID	Journal	Application
Zhaoyi Liang	36210461	Cell Mol Biol Lett	WB
Ying Ma	39235072	J Extracell Vesicles	WB
Yuling Chen	39114150	Nanoscale Adv	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67715-1-1g (Alix antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 67715-1-lg (Alix antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



WB result of Alix antibody (67715-1-Ig; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Alix transfected HEK-293 cells.



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 67715-1-Ig (Alix antibody) at dilution of 1:8000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using Alix antibody (67715-1-Ig, Clone: 1H9D9) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).