### For Research Use Only

# **GSDME** Monoclonal antibody

Catalog Number:67731-1-lg 4 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number:

67731-1-Ig BC019689
Size: GeneID (NCBI):

150ul , Concentration: 1000 ug/ml by 1687 Nanodrop and 632 ug/ml by Bradford UNIPROT ID: method using BSA as the standard; 060443

Source: Full Name:

Mouse deafness, autosomal dominant 5

Isotype: Calculated MW:
IgG1 496 aa, 55 kDa
Immunogen Catalog Number: Observed MW:
AG30514 50-55 kDa

CloneNo.: 1D9C3

**Purification Method:** 

Protein G purification

Recommended Dilutions: WB 1:1000-1:4000

**Applications** 

Tested Applications: WB, FC (Intra), ELISA Cited Applications:

WB, IHC

Species Specificity: human, mouse, rat, pig Cited Species: human, mouse, rat

#### Positive Controls:

WB: pig brain tissue, HeLa cells, HEK-293 cells, mouse brain tissue, rat brain tissue

# **Background Information**

DFNA5 (deafness, autosomal dominant 5), also known as GSDME or ICERE-1, is a 496 amino acid protein that is expressed in cochlea tissue, as well as in placenta, brain, heart, liver, lung and pancreas. Defects in the gene encoding DFNA5 are the cause of non-syndromic sensorineural deafness autosomal dominant type 5 (DFNA5), a form of sensorineural hearing loss that results from damage to one of various structures that receive sound information in the brain.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Jiewen Fu	39118110	Cancer Cell Int	WB,IHC
Wei Wang	37295545	Exp Neurol	WB
Youqing Ye	37220137	ACS Appl Mater Interfaces	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 67731-1-1g (DFNA5 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 67731-1-1g (DFNA5 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



1X10^6 SH-SY5Y cells were intracellularly stained with 0.2 ug Anti-Human DFNA5/GSDME (67731-1-1g, Clone:1D9C3) and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgC(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).