## For Research Use Only

## USP14 Monoclonal antibody

Catalog Number:67746-1-lg 5 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: Purification Method: 67746-1-lg BC003556 Protein G purification

Size:GeneID (NCBI):CloneNo.:150ul , Concentration: 1000 ug/ml by 90973B12G3

 Nanodrop;
 UNIPROT ID:
 Recommended Dilutions:

 Source:
 P54578
 WB 1:5000-1:50000

 Mouse
 Full Name:
 IF/ICC 1:200-1:800

Isotype: ubiquitin specific peptidase 14 (tRNA-

IgG1 guanine transglycosylase)

Immunogen Catalog Number: Calculated MW:

AG6312 54 kDa

Observed MW:

60 kDa

**Applications** 

Tested Applications: Positive Controls:
WB, IF/ICC, ELISA WB, HCT 116 coll.

WB; IF/ICC, ELISA

WB: HCT 116 cells, HeLa cells, Jurkat cells, HSC-T6

Cited Applications:

Cells, NIH/3T3 cells, pig brain tissue, rabbit brain

tissue, TF, IP

tissue, T16 cells, HeLa cells, Jurkat cells, HSC-T6

cells, NIH/3T3 cells, pig brain tissue, rabbit brain

tissue, rat brain tissue, mouse brain tissue, U2OS cells,

K-562 cells

IF/ICC : HepG2 cells,

Cited Species: human, mouse, rat

Species Specificity: Human, Mouse, Rat, Pig, Rabbit

**Background Information** 

USP14(Ubiquitin carboxyl-terminal hydrolase 14) is also named as TGT and belongs to the peptidase C19 family. Mammalian USP14 is unique among known UBP enzymes in that it is activated catalytically upon specific association with the 26S proteasome. USP14 inhibition accelerated the degradation of oxidized proteins and enhanced resistance to oxidative stress.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Zhiming Zhang	39490774	Free Radic Biol Med	WB
Ayako Nogami	39408703	Int J Mol Sci	WB
Haiying Rui	39178733	Redox Biol	IP,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 67746-1-1g (USP14 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using USP14 antibody (67746-1-1g, Clone: 3B12G3) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 67746-1-lg (USP14 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control