For Research Use Only

REL Monoclonal antibody

Catalog Number: 67747-1-Ig



Purification Method:

Protein G purification

Basic Information

Catalog Number: GenBank Accession Number:

67747-1-lg NM_001291746

GeneID (NCBI): CloneNo.: Size: 150ul, Concentration: 1000 ug/ml by 5966 1F6A4

Nanodrop: **UNIPROT ID:** Recommended Dilutions: Q04864 WB 1:1000-1:6000

Mouse Full Name:

Isotype: v-rel reticuloendotheliosis viral lgG1 oncogene homolog (avian)

Immunogen Catalog Number: Calculated MW:

AG26803 69 kDa

> Observed MW: 68 kDa, 80 kDa

Applications

Tested Applications:

WB, ELISA WB: Daudi cells, HEK-293 cells, Jurkat cells, MOLT-4

Positive Controls:

Species Specificity:

Human

Background Information

REL, also known as, Proto-oncogene c-Rel, is a 619 amino acid protein, which may play a role in differentiation and $lymphopoies is. \, NF-kappa-B \, is \, a \, homo- \, or \, heterodimeric \, complex \, formed \, by \, the \, Rel-like \, domain-containing \, and \, complex \, formed \, by \, the \, Rel-like \, domain-containing \, formed \, by \,$ proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52. The NF-kappa-B heterodimer RELA/p65-c-Rel is a transcriptional activator. The full-length 80-kDa c-Rel was identified (PMID: 9857058).

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67747-1-lg (REL antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.