

For Research Use Only

gp130/IL6ST Monoclonal antibody

Catalog Number: 67766-1-Ig **3 Publications**



Basic Information

Catalog Number: 67766-1-Ig	GenBank Accession Number: NM_002184	Purification Method: Protein A purification
Size: 150ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 3572	CloneNo.: 1H7C5
Source: Mouse	UNIPROT ID: P40189	Recommended Dilutions: WB 1:1000-1:4000 IHC 1:500-1:2000
Isotype: IgG1	Full Name: interleukin 6 signal transducer (gp130, oncostatin M receptor)	
Immunogen Catalog Number: AG28808	Calculated MW: 104 kDa	
	Observed MW: 130-140 kDa	

Applications

Tested Applications: WB, IHC, ELISA	Positive Controls: WB : HepG2 cells, NCCIT cells, PC-3 cells, DU 145 cells
Cited Applications: WB, IHC, IF, IP	IHC : human placenta tissue, human colon cancer tissue, human liver cancer tissue, human lung cancer tissue, human tonsillitis tissue
Species Specificity: human	
Cited Species: human	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

Glycoprotein 130 (gp130, also known as IL6ST, CD130 or IL6-beta) is a ubiquitously expressed, signal-transducing receptor that serves as the signal transduction unit for IL-6 family of cytokines, including IL-6, IL-11, IL-27, leukemia inhibitory factor (LIF), OSM, ciliary neurotrophic factor (CNTF), cardiotrophin 1 (CT-1), and cardiotrophin-like cytokine (CLC) (PMID: 9143707; 20610800). These cytokines signal through the gp130/Jak/STAT pathway (PMID: 9716487). Binding of IL-6 to IL-6R induces gp130 homodimerization and formation of a high-affinity receptor complex, which activates Jaks. That causes phosphorylation of gp130 tyrosine residues which in turn activates STAT3. gp130 is a type I transmembrane protein, and can also exist as a soluble form (sgp130). sgp130 binds to sIL-6R/IL-6 complexes and prevents their interactions with membrane-anchored gp130 on target cells (PMID: 9815830).

Notable Publications

Author	Pubmed ID	Journal	Application
Yanmiao Huo	35487760	Cell Prolif	WB,IF,IP
Yue Hu	39821173	PLoS One	WB
Yue Zhu	37306188	Cancer Med	IHC

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67766-1-Ig (gp130 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 67766-1-Ig (gp130 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 67766-1-Ig (gp130 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).