For Research Use Only

CD73 Monoclonal antibody

Catalog Number:67789-1-Ig Featured Product




Basic Information	Catalog Number:GenBank Accession Number:67789-1-lgBC015940Size:GeneID (NCBI):		lumber:	Purification Method: Protein A purification					
			CloneNo.:						
	150ul , Concentration: 1000 ug/ml by	4907		1G5G8					
	Nanodrop;	UNIPROT ID: P21589 Full Name: 5'-nucleotidase, ecto (CD73) Calculated MW:		Recommended Dilutions: WB 1:1000-1:4000 IHC 1:500-1:2000 IF-P 1:250-1:1000					
	Source:								
	Mouse								
	Isotype:								
	Immunogen Catalog Number: AG30616								
		29 kDa, 63 kDa Observed MW:							
						60-70 kDa			
	Applications	Tested Applications: WB, IHC, IF-P, ELISA Species Specificity: human, mouse, rat Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		Positive Controls: WB : HeLa cells, HepG2 cells, A549 cells, HT-29 cells, U- 87 MG cells IHC : human tonsillitis tissue, mouse kidney tissue, rat kidney tissue					
IF-P : human appendicitis tissue, human tonsillitis tissue, rat liver tissue									
Background Information						CD73, also known as ecto-5'-nucleotidase (5'-NT), is a 70-kDa, glycosyl-phosphatidylinositol-linked membrane- bound glycoprotein found in most tissues (PMID: 18404475; 20179192). CD73 is an ectoenzyme that catalyzes the dephosphorylation of AMP and other nucleoside monophosphates (PMID: 9553767). In the human immune system, CD73 is expressed on subsets of T and B cells, on germinal center follicular dendritic cells, and on thymic medullary reticular fibroblasts and epithelial cells (PMID: 2137649; 9553767). CD73 is highly expressed in many human solid tumors and is closely involved in cancer progression (PMID: 20179192).			
Storage						Storage: Store at -20°C. Stable for one year after Storage Buffer: PBS with 0.02% sodium azide and 50°	% glycerol pH 7.3.		
*** 20ul sizes contain 0.1%BSA						Aliquoting is unnecessary for -20 $^\circ$ C s	torage		


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




WB result of NT5E/CD73 antibody (67789-1-lg; 1:6000; incubated at room temperature for 1.5 hours) with sh-Control and sh-NT5E,CD73 transfected HepG2 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 67789-1-lg (NT5E/CD73 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 67789-1-1g (NT5E/CD73 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 67789-1-1g (NT5E/CD73 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human appendicitis tissue using NT5E/CD73 antibody (67789-1-1g, Clone: 1G5C8) at dilution of 1:500 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human appendicitis tissue using NT5E/CD73 antibody (67789-1-lg, Clone: 1G5C8) at dilution of 1:500 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).