For Research Use Only

CD73 Monoclonal antibody

Catalog Number:67789-1-lg Featured Product



Basic Information

Catalog Number: GenBank Accession Number:

67789-1-lg BC015940 GeneID (NCBI): Size:

150ul, Concentration: 1000 ug/ml by 4907 Nanodrop: **UNIPROT ID:** P21589 Mouse Full Name:

Isotype: 5'-nucleotidase, ecto (CD73)

lgG1 Calculated MW: Immunogen Catalog Number: 29 kDa, 63 kDa AG30616 Observed MW: 60-70 kDa

Purification Method:

Protein G purification CloneNo.:

1G5G8 Recommended Dilutions: WB 1:1000-1:4000

IHC 1:500-1:2000 IF-P 1:250-1:1000

Applications

Tested Applications: WB, IHC, IF-P, ELISA Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, HepG2 cells, A549 cells, HT-29 cells, U-87 MG cells

IHC: human tonsillitis tissue, mouse kidney tissue, rat kidney tissue

IF-P: human appendicitis tissue, human tonsillitis tissue, rat liver tissue

Background Information

CD73, also known as ecto-5'-nucleotidase (5'-NT), is a 70-kDa, glycosyl-phosphatidylinositol-linked membranebound glycoprotein found in most tissues (PMID: 18404475; 20179192). CD73 is an ectoenzyme that catalyzes the dephosphorylation of AMP and other nucleoside monophosphates (PMID: 9553767). In the human immune system, CD73 is expressed on subsets of T and B cells, on germinal center follicular dendritic cells, and on thymic medullary reticular fibroblasts and epithelial cells (PMID: 2137649; 9553767). CD73 is highly expressed in many human solid tumors and is closely involved in cancer progression (PMID: 20179192).

Storage

Store at -20°C. Stable for one year after shipment. Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


*** 20ul sizes contain 0.1% BSA

Aliquoting is unnecessary for -20°C storage


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

Selected Validation Data




WB result of NT5E/CD73 antibody (67789-1-Ig; 1:6000; incubated at room temperature for 1.5 hours) with sh-Control and sh-NT5E,CD73 transfected HepG2 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 67789-1-1g (NT5E/CD73 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 67789-1-Ig (NT5E/CD73 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 67789-1-Ig (NT5E/CD73 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human appendicitis tissue using NT5E/CD73 antibody (67789-1-1g, Clone: 1G5G8) at dilution of 1:500 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (\$A00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human appendicitis tissue using NT5E/CD73 antibody (67789-1-1g, Clone: 1G5G8) at dilution of 1:500 and Coralitie® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (\$A00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).