For Research Use Only

PKMYT1 Monoclonal antibody

Catalog Number:67806-1-lg Featured Product

2 Publications





Basic Information	Catalog Number:	GenBank Accession Number	
	67806-1-lg	NM_004203	Protein A purification
	Size:	GeneID (NCBI):	CloneNo.:
	150ul , Concentration: 1000 ug/ml by Nanodrop;		1D12D3
	Source:	UNIPROT ID: 099640	Recommended Dilutions:
	Mouse	••••	WB 1:5000-1:50000 IHC 1:500-1:2000
		Full Name:	-
	Isotype: IgG2a	protein kinase, membrane associated tyrosine/threonine 1	
	-	mmunogen Catalog Number: Calculated MW:	
	AG30849	55 kDa	
		Observed MW:	
		55 kDa	
Applications	Tested Applications:	Posi	tive Controls:
	WB, IHC, ELISA	WB	LNCaP cells, MCF-7 cells, HeLa cells, HEK-293
	Cited Applications:		s, Jurkat cells, K-562 cells, PC-12 cells, HSC-T6
	WB, IHC	cells	s, 4T1 cells, NIH/3T3 cells
	Species Specificity:	IHC	: human colon cancer tissue,
	Human, mouse, rat	inc.	
	Cited Species:		
	human, mouse		
	Note-IHC: suggested antig with TE buffer pH 9.0; (*) A	lternatively	
	antigen retrieval may be p with citrate buffer pH 6.0	erformed	
Background Information	Antigen retrieval may be p with citrate buffer pH 6.0 Protein kinase, membrane associated kinases. PKMYT1 acts as a negative re phosphorylation of Tyr14/Tyr15 and mammalian somatic cells, PKMYT1 a telophase of mitosis by inhibiting cyc	erformed d tyrosine/threonine (PKMYT) egulator of the cell cycle by in preventing cell cycle entering ffects the assembly of the Go clin activities. Abnormal exp	g into mitosis at the G2/M transition. In
	Antigen retrieval may be p with citrate buffer pH 6.0 Protein kinase, membrane associated kinases. PKMYT1 acts as a negative re phosphorylation of Tyr14/Tyr15 and p mammalian somatic cells, PKMYT1 ar telophase of mitosis by inhibiting cyo multiple types of cancers. PKMYT1 pro 32801781)	erformed d tyrosine/threonine (PKMYT) egulator of the cell cycle by in preventing cell cycle entering ffects the assembly of the Go clin activities. Abnormal exp	nactivating the CDK1-cyclinB complex through g into mitosis at the G2/M transition. In lgi apparatus and endoplasmic reticulum duri ression of PKMYT1 is closely associated with
	Antigen retrieval may be p with citrate buffer pH 6.0 Protein kinase, membrane associated kinases. PKMYT1 acts as a negative re phosphorylation of Tyr14/Tyr15 and p mammalian somatic cells, PKMYT1 ar telophase of mitosis by inhibiting cyc multiple types of cancers. PKMYT1 pro 32801781) Author Pub	erformed tyrosine/threonine (PKMYT egulator of the cell cycle by in preventing cell cycle entering ffects the assembly of the Go clin activities. Abnormal exp otein was mainly localized in	nactivating the CDK1-cyclinB complex through g into mitosis at the G2/M transition. In Igi apparatus and endoplasmic reticulum duri ression of PKMYT1 is closely associated with the cytoplasm. (PMID: 31695486, PMID: Application
	antigen retrieval may be p with citrate buffer pH 6.0Protein kinase, membrane associated kinases. PKMYT1 acts as a negative re phosphorylation of Tyr14/Tyr15 and p mammalian somatic cells, PKMYT1 at telophase of mitosis by inhibiting cyd multiple types of cancers. PKMYT1 pro 32801781)AuthorPub Kun-Huan Zhang380	erformed tyrosine/threonine (PKMYT egulator of the cell cycle by in preventing cell cycle entering ffects the assembly of the Go clin activities. Abnormal exp otein was mainly localized in med ID Journal	nactivating the CDK1-cyclinB complex throug g into mitosis at the G2/M transition. In Igi apparatus and endoplasmic reticulum duri ression of PKMYT1 is closely associated with n the cytoplasm. (PMID: 31695486, PMID: Application inh) WB
Notable Publications	antigen retrieval may be p with citrate buffer pH 6.0Protein kinase, membrane associated kinases. PKMYT1 acts as a negative re phosphorylation of Tyr14/Tyr15 and p mammalian somatic cells, PKMYT1 art telophase of mitosis by inhibiting cyd multiple types of cancers. PKMYT1 pro 32801781)AuthorPub Kun-Huan Zhang369	erformed tyrosine/threonine (PKMYT: sgulator of the cell cycle by in preventing cell cycle entering ffects the assembly of the Go clin activities. Abnormal exp otein was mainly localized in med ID Journal 14604 Adv Sci (We	nactivating the CDK1-cyclinB complex throug g into mitosis at the G2/M transition. In Igi apparatus and endoplasmic reticulum dur ression of PKMYT1 is closely associated with n the cytoplasm. (PMID: 31695486, PMID: Application inh) WB
Notable Publications	antigen retrieval may be pwith citrate buffer pH 6.0Protein kinase, membrane associatedkinases. PKMYT1 acts as a negative rephosphorylation of Tyr14/Tyr15 and jmammalian somatic cells, PKMYT1 atelophase of mitosis by inhibiting cydmultiple types of cancers. PKMYT1 pro32801781)AuthorPubKun-Huan Zhang380Linhua Yang369Storage:Storage Buffer:	erformed tyrosine/threonine (PKMYT: egulator of the cell cycle by in preventing cell cycle entering ffects the assembly of the Go clin activities. Abnormal exp otein was mainly localized in med ID Journal 14604 Adv Sci (We 88891 Cell Oncol (er shipment.	nactivating the CDK1-cyclinB complex throug g into mitosis at the G2/M transition. In Igi apparatus and endoplasmic reticulum dur ression of PKMYT1 is closely associated with n the cytoplasm. (PMID: 31695486, PMID: Application inh) WB
Notable Publications	antigen retrieval may be pwith citrate buffer pH 6.0Protein kinase, membrane associatedkinases. PKMYT1 acts as a negative rephosphorylation of Tyr14/Tyr15 and jmammalian somatic cells, PKMYT1 atelophase of mitosis by inhibiting cydmultiple types of cancers. PKMYT1 pro32801781)AuthorPubKun-Huan Zhang380Linhua Yang369Storage:Storage Buffer:PBS with 0.02% sodium azide and 50	erformed tyrosine/threonine (PKMYT: sgulator of the cell cycle by in preventing cell cycle entering ffects the assembly of the Go clin activities. Abnormal exp otein was mainly localized in med ID Journal 14604 Adv Sci (We 88891 Cell Oncol (er shipment. % glycerol pH 7.3.	nactivating the CDK1-cyclinB complex throug g into mitosis at the G2/M transition. In Igi apparatus and endoplasmic reticulum dur ression of PKMYT1 is closely associated with n the cytoplasm. (PMID: 31695486, PMID: Application inh) WB
Background Information Notable Publications Storage	antigen retrieval may be pwith citrate buffer pH 6.0Protein kinase, membrane associatedkinases. PKMYT1 acts as a negative rephosphorylation of Tyr14/Tyr15 and jmammalian somatic cells, PKMYT1 atelophase of mitosis by inhibiting cydmultiple types of cancers. PKMYT1 pro32801781)AuthorPubKun-Huan Zhang380Linhua Yang369Storage:StorageStorage Buffer:	erformed tyrosine/threonine (PKMYT: sgulator of the cell cycle by in preventing cell cycle entering ffects the assembly of the Go clin activities. Abnormal exp otein was mainly localized in med ID Journal 14604 Adv Sci (We 88891 Cell Oncol (er shipment. % glycerol pH 7.3.	nactivating the CDK1-cyclinB complex throug g into mitosis at the G2/M transition. In Igi apparatus and endoplasmic reticulum dur ression of PKMYT1 is closely associated with n the cytoplasm. (PMID: 31695486, PMID: Application inh) WB


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data







Various lysates were subjected to SDS PAGE followed by western blot with 67806-1-lg (PKMYT1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.

Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 67806-1-Ig (PKMYT 1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). WB result of PKMYT1 antibody (67806-1-1g; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PKMYT1 transfected MCF-7 cells.