

For Research Use Only

USP28 Monoclonal antibody

Catalog Number:67814-1-Ig



Basic Information

Catalog Number: 67814-1-Ig	GenBank Accession Number: BC065928	Purification Method: Protein A purification
Size: 150ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 57646	CloneNo.: 1E3D1
Source: Mouse	UNIPROT ID: Q96RU2	Recommended Dilutions: WB 1:1000-1:4000 IF/ICC 1:400-1:1600
Isotype: IgG2a	Full Name: ubiquitin specific peptidase 28	
Immunogen Catalog Number: AG11805	Calculated MW: 583aa,66 kDa; 1077aa,120 kDa	
	Observed MW: 135 kDa	

Applications

Tested Applications: WB, IF/ICC, ELISA	Positive Controls:
Species Specificity: human, mouse, rat	WB : MDA-MB-231 cells, HSC-T6 cells, HSC-T6 cells and NIH/3T3 cells., MDA-MB-468 cells, Hela cells and LNCaP cells.
	IF/ICC : HeLa cells,

Background Information

USP28 (Ubiquitin carboxyl-terminal hydrolase 28) is also named as KIAA1515. USP28 is a 53BP1-interacting protein and a putative deubiquitinating enzyme, and belongs to a family of enzymes, called deubiquitinating enzymes (DUBs), which remove ubiquitin from the ubiquitin conjugates(PMID:16901786). It can be phosphorylated upon DNA damage at Ser-67 and Ser-714, by ATM or ATR(PMID:16901786). This protein has some isoforms produced by alternative splicing with the molecular mass of 122 kDa, 119 kDa and 66 kDa, and can be detected as 130-140 kDa (PMID: 16901786).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67814-1-Ig (USP28 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control.



Various lysates were subjected to SDS PAGE followed by western blot with 67814-1-Ig (USP28 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using USP28 antibody (67814-1-Ig, Clone: 1E3D1) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).