For Research Use Only

C9orf72 Monoclonal antibody, PBS Only proteintech®

Catalog Number: 67824-1-PBS



Purification Method:

CloneNo.:

2F9A2

Protein A purification

Basic Information

Catalog Number: GenBank Accession Number:

67824-1-PBS BC020851

GeneID (NCBI): Size:

100ug, Concentration: 1mg/ml by 203228 Nanodrop; **UNIPROT ID:** Q96LT7 Source: Mouse

Isotype: chromosome 9 open reading frame 72

lgG2a Calculated MW: Immunogen Catalog Number: 481 aa, 54 kDa AG21080 Observed MW:

50 kDa

Full Name:

Applications

Tested Applications:

WB, IHC, FC (Intra), Indirect ELISA

Species Specificity: human, mouse, rat, pig


Background Information

C9ORF72 has a domain whith polymorphic hexanucleotide repeat (GGGGCC). The C9ORF72-hexanucleotide repeat expansions have been recently identified as genetic markers in amyotrophic lateral sclerosis (ALS) and frontotemporal lobar degeneration (FTLD). The C9ORF72 repeat expansions may indicate a worse prognosis in ALS. Human C9ORF72 has some isoforms with MW 54-60 kDa and 25-30 kDa. Mouse C9orf72 has some isoforms with MW 50-60 kDa and 35 kDa.


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67824-1-1g (C9orf72 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 67824-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded rat kidney tissue slide using 67824-1-lg (C90rf72 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67824-1-PBS in a different storage buffer formulation.




1X10^6 HEK-293T cells were intracellularly stained with 0.4 ug Anti-Human C9orf72 (67824-1-Ig, Clone:2F9A2) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2a Isotype Control (C1.18.4) (65208-1-Ig, Clone: C1.18.4) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 67824-1-PBS in a



Immunohistochemical analysis of paraffinembedded mouse lung tissue slide using 67824-1-Ig (C9orf72 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67824-1-PBS in a different storage buffer formulation.



Neuro-2a cells were subjected to SDS PAGE followed by western blot with 67824-1-lg (C9orf72 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67824-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse intestine tissue slide using 67824-1-lg (C9orf72 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67824-1-PBS in a different storage buffer formulation