For Research Use Only

BAD Monoclonal antibody

Catalog Number: 67830-1-lg 8 Publications



Basic Information

Catalog Number: GenBank Accession Number:

67830-1-lg BC001901 Protein A purification

Size:GeneID (NCBI):CloneNo.:150ul , Concentration: 1000 ug/ml by 5724B12G2

 Nanodrop;
 UNIPROT ID:
 Recommended Dilutions:

 Source:
 Q92934
 WB 1:1000-1:6000

 Mouse
 Full Name:
 IF/ICC 1:200-1:800

Isotype: BCL2-associated agonist of cell death

IgG1 Calculated MW:

18 kDa Observed MW: 18-25 kDa

Applications

Tested Applications: WB, IF/ICC, ELISA Cited Applications:

WB

Species Specificity: Human, Mouse, Rat Cited Species: human, mouse, rat Positive Controls:

WB: LNCaP cells, RAW 264.7 cells, HeLa cells, HEK-293 cells, Jurkat cells, Raji cells, THP-1 cells, HSC-T6 cells

Purification Method:

IF/ICC: HeLa cells,

Background Information

BAD, also named as BBC6 and BCL2L8, promotes cell death. It is successfully competes for the binding to Bcl-X(L), Bcl-2 and Bcl-W, thereby affecting the level of heterodimerization of these proteins with BAX. BAD can reverse the death repressor activity of Bcl-X(L), but not that of Bcl-2. It appears to act as a link between growth factor receptor signaling and the apoptotic pathways.

Notable Publications

Author	Pubmed ID	Journal	Application
Yaling Zhang	36062194	Oxid Med Cell Longev	WB
Xueying Chen	33817830	FASEBJ	WB
Can Wang	39939853	J Nanobiotechnology	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67830-1-lg (BAD antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



Immunofluorescent analysis of (-20°C Methanol) fixed Hela cells using BAD antibody (67830-1-lg, Clone: 4B12G2) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L).