## For Research Use Only

## HNRNPA1 Monoclonal antibody

Catalog Number:67844-1-lg 2 Publications



**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** Protein A purification

67844-1-lg Size:

GeneID (NCBI):

150ul, Concentration: 500 ug/ml by

CloneNo.:

Nanodrop:

**UNIPROT ID:** 

BC009600

1A5A4

Mouse

P09651 Full Name: Recommended Dilutions: WB 1:20000-1:100000

Isotype: lgG2b

AG17415

heterogeneous nuclear ribonucleoprotein A1

IHC 1:2500-1:10000 IF/ICC 1:200-1:800

Immunogen Catalog Number:

Calculated MW:

34 kDa

Observed MW: 34 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, ELISA

**Cited Applications:** 

WB, IHC

Species Specificity:

human, mouse

**Cited Species:** human, mouse Positive Controls:

WB: LNCaP cells, NIH/3T3 cells, RAW 264.7 cells, HeLa

cells, Jurkat cells, K-562 cells, HEK-293 cells

IHC: human lung cancer tissue, human breast cancer tissue, human colon cancer tissue, human liver cancer

tissue

IF/ICC : HeLa cells.

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

**Background Information** 

In eukaryotic cells, nascent RNA polymerase II transcripts are associated in the nucleus with specific proteins to form ribonucleoprotein complexes called HNRP or 40S. Protein moiety of the HNRP particle has 6 major components called core proteins, and HNRPA1 is one of them. hnRNP A1 is a multifunctional RNA binding protein which is involved in pre-mRNA splicing, export of mature transcripts from the nucleus, mRNA turnover, and internal ribosome entry site (IRES)-mediated translation. HNRNPA1 has three isoforms with MW 30 kDa, 34 kDa and 39 kDa.

**Notable Publications** 

Author	Pubmed ID	Journal	Application
Xiang Ma	39428563	CNS Neurosci Ther	WB
Cheng Lian	39341825	Signal Transduct Target Ther	WB,IHC

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 67844-1-1g (HNRNPA1 antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.




Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 67844-1-Ig (HNRNPA1 antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 67844-1-1g (HNRNPA1 antibody) at dilution of 1:5000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using HNRNPA1 antibody (67844-1-lg, Clone: 1A5A4) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM002), CL594-phalloidin (red).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 67844-1-1g (HNRNPA1 antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).