For Research Use Only

CDK7 Monoclonal antibody

Catalog Number: 67889-1-Ig



Basic Information

Catalog Number: GenBank Accession Number:

67889-1-lg BC005298 Size: GeneID (NCBI):

150ul , Concentration: 1000 $\mu g/ml$ by 1022 1E7B6 Nanodrop;

Source: cyclin-dependent kinase 7 Mouse Calculated MW: Isotype: 39 kDa

lgG1 Observed MW: Immunogen Catalog Number: 39 kDa

AG25737

Applications

Tested Applications:

IF, IHC, WB, ELISA Species Specificity: Human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method: Protein G purification

CloneNo.:

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:2000 IF 1:750-1:3000

Positive Controls:

WB: A549 cells, U2OS cells, MCF-7 cells, Hela cells, Jurkat cells, HEK-293 cells, K-562 cells, HSC-T6 cells,

NIH/3T3 cells, 4T1 cells

IHC: human breast cancer tissue, IF: HeLa cells, A431 cells

Background Information

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67889-1-lg (CDK7 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 67889-1-Ig (CDK7 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 67889-1-Ig (CDK7 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CDK7 antibody (67889-1-1g, Clone: 1E7B6) at dilution of 1:1500 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed A431 cells using CDK7 antibody (67889-1-Ig, Clone: 1E7B6) at dilution of 1:1500 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 67889-1-1g (CDK7 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.