

For Research Use Only

HSPB3 Monoclonal antibody

Catalog Number: 67890-1-Ig



Basic Information

Catalog Number: 67890-1-Ig	GenBank Accession Number: BC126272	Purification Method: Protein A purification
Size: 150ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 8988	CloneNo.: 2A10B5
Source: Mouse	UNIPROT ID: Q12988	Recommended Dilutions: WB: 1:5000-1:50000 IHC: 1:250-1:1000 IF/ICC: 1:400-1:1600 FC (Intra): 0.50 ug per 10 ⁶ cells in a 100 µl suspension
Isotype: IgG2b	Full Name: heat shock 27kDa protein 3	
Immunogen Catalog Number: AG26940	Calculated MW: 150 aa, 17 kDa	
	Observed MW: 20 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, FC (Intra), ELISA	Positive Controls:
Species Specificity: human, mouse, rat, rabbit	WB : human heart tissue, mouse heart tissue, rabbit heart tissue, rat heart tissue
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IHC : rat kidney tissue, mouse kidney tissue
	IF/ICC : HeLa cells,
	FC (Intra) : HeLa cells,

Background Information

Heat shock protein family B (small) member 3 (HSPB3) is the third member of the sHSP family in human and is mainly expressed in skeletal and smooth muscles, and can associate with HSPB2, which is another sHSP prominently expressed in cardiac and skeletal muscles. HSPB3 is involved in several cellular functions such as cell signaling, differentiation, and apoptosis, and mutations of HSPB3 are associated with neuromuscular diseases, such as axonal Charcot-Marie-Tooth disease (PMID: 31709619) (PMID: 32093037). The molecular weight of HSPB3 is about 17 kDa.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67890-1-Ig (HSPB3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.




human heart tissue were subjected to SDS PAGE followed by western blot with 67890-1-Ig (HSPB3 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 67890-1-Ig (HSPB3 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 67890-1-Ig (HSPB3 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 67890-1-Ig (HSPB3 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded rat kidney tissue slide using 67890-1-Ig (HSPB3 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded rat kidney tissue slide using 67890-1-Ig (HSPB3 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using HSPB3 antibody (67890-1-Ig, Clone: 2A10B5) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using HSPB3 antibody (67890-1-Ig, Clone: 2A10B5) at dilution of 1:800 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L).



1X10⁶ HeLa cells were intracellularly stained with 0.5 ug Anti-Human HSPB3 (67890-1-Ig, Clone:2A10B5) and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.5 ug Mouse IgG2b Isotype Control (66360-3-Ig, Clone: K11B8C4B5) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).