

For Research Use Only

HSPB3 Monoclonal antibody, PBS Only



Catalog Number: 67890-1-PBS

Basic Information

Catalog Number: 67890-1-PBS	GenBank Accession Number: BC126272	Purification Method: Protein A purification
Size: 100ug , Concentration: 1mg/ml by Nanodrop;	GeneID (NCBI): 8988	CloneNo.: 2A10B5
Source: Mouse	UNIPROT ID: Q12988	
Isotype: IgG2b	Full Name: heat shock 27kDa protein 3	
Immunogen Catalog Number: AG26940	Calculated MW: 150 aa, 17 kDa	
	Observed MW: 20 kDa	

Applications

Tested Applications:
Indirect ELISA, IF/ICC, IHC, WB

Species Specificity:
rabbit, rat, mouse, human

Background Information

Heat shock protein family B (small) member 3 (HSPB3) is the third member of the sHSP family in human and is mainly expressed in skeletal and smooth muscles, and can associate with HSPB2, which is another sHSP prominently expressed in cardiac and skeletal muscles. HSPB3 is involved in several cellular functions such as cell signaling, differentiation, and apoptosis, and mutations of HSPB3 are associated with neuromuscular diseases, such as axonal Charcot-Marie-Tooth disease (PMID: 31709619) (PMID: 32093037). The molecular weight of HSPB3 is about 17 kDa.

Storage


Storage:
Store at -80°C.

Storage Buffer:
PBS Only


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67890-1-Ig (HSPB3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 67890-1-PBS in a different storage buffer formulation.




human heart tissue were subjected to SDS PAGE followed by western blot with 67890-1-Ig (HSPB3 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67890-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded rat kidney tissue slide using 67890-1-Ig (HSPB3 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67890-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 67890-1-Ig (HSPB3 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67890-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 67890-1-Ig (HSPB3 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67890-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 67890-1-Ig (HSPB3 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67890-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded rat kidney tissue slide using 67890-1-Ig (HSPB3 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67890-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using HSPB3 antibody (67890-1-Ig, Clone: 2A10B5) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67890-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using HSPB3 antibody (67890-1-Ig, Clone: 2A10B5) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67890-1-PBS in a different storage buffer formulation.



1X10⁶ HeLa cells were intracellularly stained with 0.5 ug Anti-Human HSPB3 (67890-1-Ig, Clone:2A10B5) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.5 ug Mouse IgG2b Isotype Control (66360-3-Ig, Clone: K11B8C4B5) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 67890-1-PBS in a

