

For Research Use Only

Stomatin Monoclonal antibody

Catalog Number: 67894-1-Ig



Basic Information

Catalog Number: 67894-1-Ig	GenBank Accession Number: BC010703	Purification Method: Protein G purification
Size: 150ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 2040	CloneNo.: 1A4D6
Source: Mouse	Full Name: stomatin	Recommended Dilutions: WB 1:2000-1:10000 IHC 1:50-1:500
Isotype: IgG1	Calculated MW: 288 aa, 32 kDa	
Immunogen Catalog Number: AG2676	Observed MW: 32 kDa	

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

Human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : human placenta tissue, HeLa cells, human peripheral blood platelets, human testis tissue

IHC : human lung cancer tissue,

Background Information

STOM, also named as BND7, EPB72, Stomatin and Protein 7.2b, belongs to the band 7/mec-2 family. It is thought to regulate cation conductance. STOM may regulate ACCN1 and ACCN3 gating. STOM is a membrane protein involved in regulation of monovalent cation transport through lipid membranes. Interestingly, STOM (which has a structure similar to caveolin) is a major lipid-raft component of erythrocytes. It may have a role in membrane microdomains modulation leading to membrane budding and MPs release.(PMID:19138258)

Storage

Storage:

Store at -20°C.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



human placenta tissue were subjected to SDS PAGE followed by western blot with 67894-1-Ig (Stomatin antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 67894-1-Ig (Stomatin antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).