For Research Use Only

RAB11A Monoclonal antibody, PBS Only proteintech

Catalog Number:67902-1-PBS

## Basic Information Catalog Number: 67902-1-PBS Size: 100ug , Concentration: 1mg, Nanodrop;

67902-1-PBS Size: 100ug , Concentration: 1mg/ml by Nanodrop; Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG8879 GenBank Accession Number: BC013348 GeneID (NCBI): 8766 UNIPROT ID: P62491 Full Name: RAB11A, member RAS oncogene family Calculated MW: 216 aa, 24 kDa Observed MW: 24 kDa Purification Method: Protein A purification CloneNo.: 4A4C9

Antibodies | ELISA kits | Proteins

www.ptglab.com

## **Applications**

Tested Applications: WB, IF/ICC, FC (Intra), ELISA Species Specificity: human, mouse, rat, pig, rabbit

**Background Information** 

Rab11a, Rab11b and Rab25 comprise a subfamily of Rab proteins that belong to the Ras-related small GTPase protein superfamily (PMID: 8074219, PMID: 10708602). Rab11 proteins are involved in endocytosis and plasma membrane recycling in non-polarized and polarized cells (PMID: 8638719, PMID: 8922376). Rab11a and Rab11b are ubiquitously expressed in most tissues, while Rab25 is only expressed in epithelial cells. Rab11a is known to regulate the recycling of internalized cell surface proteins and receptors from the early endosome through the trans-Golgi network (PMID: 11121436, PMID: 9600939).


## Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67902-1-lg (RAB11A antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 67902-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 67902-1-lg (RAB11A antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 67902-1-PBS in a different storage buffer formulation.



mouse brain tissue were subjected to SDS PAGE followed by western blot with 67902-1-Ig (RAB11A antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67902-1-PBS in a different storage buffer formulation.



1x10^6 HeLa cells were intracellularly stained with 0.4 ug RAB11A Monoclonal antibody (67902-1-lg, Clone:4A4C9) and CoraLite@488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1)(red), or 0.4 ug Mouse IgG1 isotype control Mouse McAb (66360-1-lg, Clone: 1F8D3) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 67902-1-PBS in a different storage buffer



Immunofluorescent analysis of (-20°C Methanol) fixed A549 cells using RAB11A antibody (67902-1-Ig, Clone: 4A4C9) at dilution of 1:800 and Coralite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1). This data was developed using the same antibody clone with 67902-1-PBS in a different storage buffer formulation.