For Research Use Only

YTHDC1 Monoclonal antibody

Catalog Number:67911-1-lg 1 Publications



Basic Information

Catalog Number: GenBank Accession Number:

67911-1-Ig BC053863

Size: GeneID (NCBI): 150ul , Concentration: 1000 ug/ml by 91746

Nanodrop; UNIPROT ID:
Source: Q96MU7
Mouse Full Name:

Isotype: YTH domain containing 1

IgG1 Calculated MW:
Immunogen Catalog Number: 85 kDa

AG6289 Observed MW: ~100 kDa

Applications

Tested Applications: WB, IHC, ELISA Cited Applications:

WB

Species Specificity: Human, mouse Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: LNCaP cells, MDA-MB-231 cells, HeLa cells, HEK-

Purification Method:

Protein G purification

Recommended Dilutions:

WB 1:2000-1:10000

IHC 1:2000-1:8000

CloneNo.:

1F5G11

293 cells, Jurkat cells, K-562 cells

IHC: human breast cancer tissue, human cervical

cancer tissue, mouse testis tissue

Background Information

YTHDC1, containing 1 YTH domain, is one of the m6A-binding proteins that can recognize and bind to the m6A methylation site and plays a specific role in gene expression. YTHDC1 is constitutively enriched in the nucleus. It regulates mRNA splicing by bridging the interactions between the trans- and cis-regulatory elements to bind targeted mRNAs. YTHDC1 mediates the export of m6A modified mRNA from the nucleus to the cytoplasm by interacting with SRSF3, which is an essential factor driving the tumorigenic process in various types of cancers including breast cancer, colon cancer, ovarian cancer, osteosarcoma, and glioblastoma (32368386).

Notable Publications

Author	Pubmed ID	Journal	Application
Xiaoxin Xu	39092767	Mol Carcinog	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67911-1-1g (YTHDC1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with Lamin B1 Monoclonal antibody (66095-1-1g) as loading control.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 67911-1-1g (YTHDC1 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 67911-1-1g (YTHDC1 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 67911-1-1g (YTHDC1 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 67911-1-Ig (YTHDC 1 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 67911-1-Ig (YTHDC 1 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).