

For Research Use Only

# YTHDC1 Monoclonal antibody

Catalog Number: 67911-1-Ig



## Basic Information

|  |  |   |
|--|--|---|
| <b>Catalog Number:</b><br>67911-1-Ig                           | <b>GenBank Accession Number:</b><br>BC053863 | <b>Purification Method:</b><br>Protein G purification                   |
| <b>Size:</b><br>150ul , Concentration: 1000 µg/ml by Nanodrop; | <b>GeneID (NCBI):</b><br>91746               | <b>CloneNo.:</b><br>1F5G11  |
| <b>Source:</b><br>Mouse  | <b>Full Name:</b><br>YTH domain containing 1 | <b>Recommended Dilutions:</b><br>WB 1:2000-1:10000<br>IHC 1:2000-1:8000 |
| <b>Isotype:</b><br>IgG1  | <b>Calculated MW:</b><br>85 kDa              |   |
| <b>Immunogen Catalog Number:</b><br>AG6289                     | <b>Observed MW:</b><br>~100 kDa              |   |

## Applications

### Tested Applications:

IHC, WB, ELISA

### Species Specificity:

Human, mouse

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

**WB** : LNCaP cells, MDA-MB-231 cells, HeLa cells, HEK-293 cells, Jurkat cells, K-562 cells

**IHC** : human breast cancer tissue, human cervical cancer tissue, mouse testis tissue

## Background Information

YTHDC1, containing 1 YTH domain, is one of the m6A-binding proteins that can recognize and bind to the m6A methylation site and plays a specific role in gene expression. YTHDC1 is constitutively enriched in the nucleus. It regulates mRNA splicing by bridging the interactions between the trans- and cis-regulatory elements to bind targeted mRNAs. YTHDC1 mediates the export of m6A modified mRNA from the nucleus to the cytoplasm by interacting with SRSF3, which is an essential factor driving the tumorigenic process in various types of cancers including breast cancer, colon cancer, ovarian cancer, osteosarcoma, and glioblastoma (32368386).

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67911-1-Ig (YTHDC1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with Lamin B1 Monoclonal antibody (66095-1-Ig) as loading control.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 67911-1-Ig (YTHDC1 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 67911-1-Ig (YTHDC1 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 67911-1-Ig (YTHDC1 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 67911-1-Ig (YTHDC1 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 67911-1-Ig (YTHDC1 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).