### For Research Use Only

# USP10 Monoclonal antibody

Catalog Number:67917-1-lg

Featured Product

1 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number:

67917-1-lg BC000263 Protein A purification Size: GeneID (NCBI): CloneNo.:

150ul , Concentration: 1000 ug/ml by 9100 1B10C9
Nanodrop; UNIPROT ID: Recomm

Source: Q14694 Mouse Full Name:

Isotype: ubiquitin specific peptidase 10

IgG1 Calculated MW:
Immunogen Catalog Number: 87 kDa, 92 kDa
AG7233 Observed MW:
100-110 kDa

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:2000 IF/ICC 1:200-1:800

**Purification Method:** 

**Applications** 

Tested Applications: WB, IHC, IF/ICC, ELISA

Cited Applications:

WB

Species Specificity: Human, mouse, rat Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: LNCaP cells, MCF-7 cells, HeLa cells, HEK-293 cells, Jurkat cells, K-562 cells, HSC-T6 cells, PC-12

cells, NIH/3T3 cells

IHC: human liver cancer tissue,

IF/ICC: HEK-293 cells,

## **Background Information**

USP10, a cytoplasmic ubiquitin-specific protease, deubiquitinates p53, reversing Mdm2-induced p53 nuclear export and degradation. After DNA damage, USP10 is stabilized, and a fraction of USP10 translocates to the nucleus to activate p53. The translocation and stabilization of USP10 is regulated by ATM -mediated phosphorylation of USP10 at Thr42 and Ser337. (PMID: 20096447, PMID: 26876938)

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Xiaoli Wang	38363675	Cell Rep	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


### **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 67917-1-1g (USP10 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67917-1-Ig (USP10 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using USP10 antibody (67917-1-lg, Clone: 1B10C9) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L).



WB result of USP10 antibody (67917-1-lg; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-USP10 transfected MCF-7 cells.