For Research Use Only

## ValRS Monoclonal antibody

Catalog Number:67935-1-lg





## Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** 67935-1-lg BC012808 Protein A purification GenelD (NCBI): CloneNo.: Size: 150ul , Concentration: 1000 ug/ml by 7407 1A2C3 Nanodrop: UNIPROT ID: Recommended Dilutions: Source: P26640 WB 1:5000-1:50000 Mouse IHC 1:250-1:1000 Full Name: IF/ICC 1:400-1:1600 Isotype: valyl-tRNA synthetase lgG2a Calculated MW: Immunogen Catalog Number: 1264 aa, 140 kDa AG9312 **Observed MW:** 140 kDa **Applications Tested Applications:** Positive Controls: WB, IHC, IF/ICC, ELISA WB: HEK-293 cells, LNCaP cells, MCF-7 cells, HepG2 Species Specificity: cells, HeLa cells, K-562 cells, HSC-T6 cells, NIH/3T3 Human, mouse, rat cells, 4T1 cells Note-IHC: suggested antigen retrieval with IHC : human tonsillitis tissue. TE buffer pH 9.0; (\*) Alternatively, antigen IF/ICC : HeLa cells, retrieval may be performed with citrate buffer pH 6.0 Storage: Storage Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage \*\*\* 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using ValRS antibody (67935-1-lg, Clone: 1A2C3) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 67935-1-1g (VaIRS antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Various lysates were subjected to SDS PAGE followed by western blot with 67935-1-lg (VaIRS antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with Lamin B1 Monoclonal antibody (66095-1-lg) as loading control.