For Research Use Only

HADHB Monoclonal antibody

Catalog Number: 67967-1-Ig



Purification Method:

IF/ICC 1:200-1:800

CloneNo.:

Protein A purification

Basic Information

Catalog Number: GenBank Accession Number:

67967-1-lg BC017564
Size: GeneID (NCBI):

 150ul , Concentration: 1000 ug/ml by 3032
 1D12F4

 Nanodrop;
 UNIPROT ID:
 Recommended Dilutions:

 Source:
 P55084
 WB 1:5000-1:50000

 Mouse
 Full Name:
 IHC 1:1000-1:4000

 Isotype:
 hydroxyacyl-Coenzyme A

 IgG2b
 dehydrogenase/3-ketoacyl

 Immunogen Catalog Number:
 Coenzyme A thiolase/enoyl

AG30315 Coenzyme A hydratase (trifunctional

protein), beta subunit Calculated MW:

Observed MW: 47 kDa

51 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Species Specificity:

human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: LNCaP cells, pig heart tissue, Hela cells, HEK-293 cells, HepG2 cells, Jurkat cells, Rabbit heart tissues, Rat heart tissues, Mouse heart tissues

IHC: human colon cancer tissue,

IF/ICC : HeLa cells,

Background Information

HADHB, also named as TP- beta, Acetyl-CoA acyltransferase and Beta-ketothiolase, is a mitochondrial trifunctional enzyme subunit beta. Mitochondrial trifunctional enzyme catalyzes the last three of the four reactions of the mitochondrial beta-oxidation pathway. The mitochondrial beta-oxidation pathway is the major energy-producing process in tissues and is performed through four consecutive reactions breaking down fatty acids into acetyl-CoA. Among the enzymes involved in this pathway, the trifunctional enzyme exhibits specificity for long-chain fatty acids. Mitochondrial trifunctional enzyme is a heterotetrameric complex composed of two proteins, the trifunctional enzyme subunit alpha/HADHA carries the 2,3-enoyl-CoA hydratase and the 3-hydroxyacyl-CoA dehydrogenase activities, while the trifunctional enzyme subunit beta/HADHB described here bears the 3-ketoacyl-CoA thiolase activity. HADHB has 2 isoforms produced by alternative splicing with the MW of 49 kDa and 51 kDa.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffe

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67967-1-lg (HADHB antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control



Various lysates were subjected to SDS PAGE followed by western blot with 67967-1-1g (HADHB antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 67967-1-Ig (HADHB antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using HADHB antibody (67967-1-1g, Clone: 1D12F4) at dilution of 1:400 and CoraLite®488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 67967-1-lg (HADHB antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using HADHB antibody (67967-1-1g, Clone: 1D12F4) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).