

For Research Use Only

HADHB Monoclonal antibody

Catalog Number: 67967-1-Ig



Basic Information

Catalog Number: 67967-1-Ig	GenBank Accession Number: BC017564	Purification Method: Protein A purification
Size: 150ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 3032	CloneNo.: 1D12F4
Source: Mouse	Full Name: hydroxyacyl-Coenzyme A dehydrogenase/3-ketoacyl- Coenzyme A thiolase/enoyl- Coenzyme A hydratase (trifunctional protein), beta subunit	Recommended Dilutions: WB 1:5000-1:50000 IF 1:200-1:800
Isotype: IgG2b	Calculated MW: 51 kDa	
Immunogen Catalog Number: AG30315	Observed MW: 47 kDa	

Applications

Tested Applications:

IF, WB, ELISA

Species Specificity:

Human, Mouse, Rat, Pig

Positive Controls:

WB : LNCaP cells, pig heart tissue, HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, Rabbit heart tissues, Rat heart tissues, Mouse heart tissues

IF : HeLa cells,

Background Information

HADHB, also named as TP- beta, Acetyl-CoA acyltransferase and Beta-ketothiolase, is a mitochondrial trifunctional enzyme subunit beta. Mitochondrial trifunctional enzyme catalyzes the last three of the four reactions of the mitochondrial beta-oxidation pathway. The mitochondrial beta-oxidation pathway is the major energy-producing process in tissues and is performed through four consecutive reactions breaking down fatty acids into acetyl-CoA. Among the enzymes involved in this pathway, the trifunctional enzyme exhibits specificity for long-chain fatty acids. Mitochondrial trifunctional enzyme is a heterotetrameric complex composed of two proteins, the trifunctional enzyme subunit alpha/HADHA carries the 2,3-enoyl-CoA hydratase and the 3-hydroxyacyl-CoA dehydrogenase activities, while the trifunctional enzyme subunit beta/HADHB described here bears the 3-ketoacyl-CoA thiolase activity. HADHB has 2 isoforms produced by alternative splicing with the MW of 49 kDa and 51 kDa.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67967-1-Ig (HADHB antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.




Various lysates were subjected to SDS PAGE followed by western blot with 67967-1-Ig (HADHB antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 67967-1-Ig (HADHB antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using HADHB antibody (67967-1-Ig, Clone: 1D12F4) at dilution of 1:400 and Coralite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using HADHB antibody (67967-1-Ig, Clone: 1D12F4) at dilution of 1:800 and Coralite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).