For Research Use Only

## PRMT6 Monoclonal antibody

Catalog Number:67981-1-lg 1 Publications



Basic Information	Catalog Number: 67981-1-lg	GenBank Accession Number: BC002729	Purification Method: Protein A purification
	Size:	GeneID (NCBI):	CloneNo.:
	150ul , Concentration: 1000 ug/ml by Nanodrop; Source:	55170	4B5A9
		UNIPROT ID:	Recommended Dilutions:
	Mouse	Q96LA8 Full Name:	WB 1:5000-1:50000
	Isotype:	protein arginine methyltransferase 6 Calculated MW:	
	lgG2a		
	Immunogen Catalog Number: AG7934	42 kDa	
		Observed MW:	
		42 kDa	
Applications	Tested Applications:	Positive Controls:	
	WB, ELISA	WB : HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, K-562 cells, HSC-T6 cells, NIH/3T3 cells	
	Cited Applications: WB		
	Species Specificity: Human, Mouse, Rat		
	Cited Species: human		
	Protein arginine methyltransferase 6 (PRMT6) is a type I PRMT which is involved in epigenetic regulation of gene expression, alternative splicing, development and differentiation, DNA repair, cell proliferation and senescence, DNA methylation, mitosis, inflammation, innate antiviral immunity, spermatogenesis, transactivation of nuclear receptors and cell signaling. The human PRMT6 gene, located on Chromosome 1, encodes for the 41.9 kDa PRMT6 enzyme. PRMT6 is predominantly localized to the nucleus, in stark contrast to PRMT3 and PRMT5 which are preponderantly cytosolic, while other PRMTs are found in both nucleus and cytosol. PRMT6 is expressed in a wide range of tissues with high expression in kidney and testes. PRMT6 generates asymmetric dimethylation modifications in histone 3 at arginine 2, arginine 17 and arginine 42 (H3R2me2a, H3R17me2a and H3R42me2a) an in histone H2A at arginine 26 (H2AR26me2a) and participates in the epigenetic regulation of gene expression.		
Background Information	expression, alternative splicing, deve DNA methylation, mitosis, inflamma receptors and cell signaling. The hum enzyme. PRMT6 is predominantly loc preponderantly cytosolic, while other range of tissues with high expression modifications in histone 3 at argining	elopment and differentiation, DNA tion, innate antiviral immunity, sp an PRMT6 gene, located on Chrom alized to the nucleus, in stark contu PRMTs are found in both nucleus a in kidney and testes. PRMT6 gene a 2, arginine 17 and arginine 42 (H2	repair, cell proliferation and senescence, ermatogenesis, transactivation of nuclea osome 1, encodes for the 41.9 kDa PRMT6 rast to PRMT3 and PRMT5 which are and cytosol. PRMT6 is expressed in a wide rates asymmetric dimethylation 3R2me2a, H3R17me2a and H3R42me2a) a
Background Information	expression, alternative splicing, deve DNA methylation, mitosis, inflamma receptors and cell signaling. The hum enzyme. PRMT6 is predominantly loc preponderantly cytosolic, while other range of tissues with high expression modifications in histone 3 at arginine in histone H2A at arginine 26 (H2AR2	elopment and differentiation, DNA tion, innate antiviral immunity, sp an PRMT6 gene, located on Chrom alized to the nucleus, in stark contu PRMTs are found in both nucleus a in kidney and testes. PRMT6 gene a 2, arginine 17 and arginine 42 (H2	repair, cell proliferation and senescence, ermatogenesis, transactivation of nuclear osome 1, encodes for the 41.9 kDa PRMT6 rast to PRMT3 and PRMT5 which are and cytosol. PRMT6 is expressed in a wide rates asymmetric dimethylation 3R2me2a, H3R17me2a and H3R42me2a) a
	expression, alternative splicing, deve DNA methylation, mitosis, inflamma receptors and cell signaling. The hum enzyme. PRMT6 is predominantly loc preponderantly cytosolic, while other range of tissues with high expression modifications in histone 3 at arginine in histone H2A at arginine 26 (H2AR2 Author Pub	elopment and differentiation, DNA tion, innate antiviral immunity, sp an PRMT6 gene, located on Chrom alized to the nucleus, in stark conti PRMTs are found in both nucleus a in kidney and testes. PRMT6 gene 2, arginine 17 and arginine 42 (H 6me2a) and participates in the ep	repair, cell proliferation and senescence, ermatogenesis, transactivation of nuclear osome 1, encodes for the 41.9 kDa PRMT6 rast to PRMT3 and PRMT5 which are and cytosol. PRMT6 is expressed in a wide rates asymmetric dimethylation 3R2me2a, H3R17me2a and H3R42me2a) a genetic regulation of gene expression.
	expression, alternative splicing, deve DNA methylation, mitosis, inflamma receptors and cell signaling. The hum enzyme. PRMT6 is predominantly loc preponderantly cytosolic, while other range of tissues with high expression modifications in histone 3 at arginine in histone H2A at arginine 26 (H2AR2 Author Pub	Peopment and differentiation, DNA tion, innate antiviral immunity, sp an PRMT6 gene, located on Chrom alized to the nucleus, in stark contr PRMTs are found in both nucleus a in kidney and testes. PRMT6 gene e 2, arginine 17 and arginine 42 (H2 6me2a) and participates in the epi med ID Journal	repair, cell proliferation and senescence, ermatogenesis, transactivation of nuclear osome 1, encodes for the 41.9 kDa PRMT6 rast to PRMT3 and PRMT5 which are and cytosol. PRMT6 is expressed in a wide rates asymmetric dimethylation 3R2me2a, H3R17me2a and H3R42me2a) a genetic regulation of gene expression.
Notable Publications	expression, alternative splicing, deve DNA methylation, mitosis, inflamma receptors and cell signaling. The hum enzyme. PRMT6 is predominantly loc preponderantly cytosolic, while other range of tissues with high expression modifications in histone 3 at arginine in histone H2A at arginine 26 (H2AR2 Author Pub	Pelopment and differentiation, DNA tion, innate antiviral immunity, sp lan PRMT6 gene, located on Chrom alized to the nucleus, in stark conti PRMTs are found in both nucleus a in kidney and testes. PRMT6 gene e 2, arginine 17 and arginine 42 (H 6me2a) and participates in the epi med ID Journal 24136 Leukemia	repair, cell proliferation and senescence, ermatogenesis, transactivation of nuclea osome 1, encodes for the 41.9 kDa PRMT6 rast to PRMT3 and PRMT5 which are and cytosol. PRMT6 is expressed in a wide rates asymmetric dimethylation 3R2me2a, H3R17me2a and H3R42me2a) a genetic regulation of gene expression.
	expression, alternative splicing, deve DNA methylation, mitosis, inflamma receptors and cell signaling. The hum enzyme. PRMT6 is predominantly loc preponderantly cytosolic, while other range of tissues with high expression modifications in histone 3 at arginine in histone H2A at arginine 26 (H2AR2 Author Pub Xinyu Yang 384 Storage: Store at -20°C. Stable for one year aft Storage Buffer:	Pelopment and differentiation, DNA tion, innate antiviral immunity, sp lan PRMT6 gene, located on Chrom alized to the nucleus, in stark conti PRMTs are found in both nucleus a in kidney and testes. PRMT6 gene e 2, arginine 17 and arginine 42 (H 6me2a) and participates in the epi med ID Journal 24136 Leukemia er shipment. % glycerol pH 7.3.	repair, cell proliferation and senescence, ermatogenesis, transactivation of nuclea osome 1, encodes for the 41.9 kDa PRMT6 rast to PRMT3 and PRMT5 which are and cytosol. PRMT6 is expressed in a wide rates asymmetric dimethylation 3R2me2a, H3R17me2a and H3R42me2a) a genetic regulation of gene expression.

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67981-1-Ig (PRMT6 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.