For Research Use Only

ACAT2 Monoclonal antibody

Catalog Number:68005-1-lg Featured Product

1 Publications



Basic Information

Catalog Number: GenBank Accession Number:

68005-1-lg BC000408 Protein G purification

GeneID (NCBI): CloneNo.: 150ul, Concentration: 1000 ug/ml by 39 1F5F11

Nanodrop: **UNIPROT ID:** Recommended Dilutions: Q9BWD1 WB 1:5000-1:50000 Mouse IHC 1:200-1:800 Full Name: acetyl-Coenzyme A acetyltransferase IF/ICC 1:500-1:2000 Isotype:

lgG1

Immunogen Catalog Number: Calculated MW: 41 kDa AG6596

> Observed MW: 42 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA Cited Applications:

Species Specificity: Human, mouse, rat, rabbit

Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HepG2 cells, HCT 116 cells, rat brain tissue, HeLa cells, Jurkat cells, K-562 cells, mouse brain tissue,

Purification Method:

rabbit brain tissue

IHC: human placenta tissue, IF/ICC: HepG2 cells,

Notable Publications

Author Pubmed ID Journal Application Yang Wang 39059092 Phytomedicine WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Various cell lysates were subjected to SDS PAGE followed by western blot with 68005-1-lg (ACAT2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 68005-1-Ig (ACAT2 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using ACAT2 antibody (68005-1-lg, Clone: 1F5F11) at dilution of 1:1000 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L).



WB result of ACAT2 antibody (68005-1-Ig; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ACAT2 transfected HepG2 cells.