For Research Use Only

STK11/LKB1 Monoclonal antibody

Catalog Number: 68016-1-Ig



Basic Information

Catalog Number: GenBank Accession Number:

68016-1-lg BC007981 GeneID (NCBI): 150ul , Concentration: 500 µg/ml by

Nanodrop: **UNIPROT ID:** Q15831 Mouse Full Name:

Isotype: serine/threonine kinase 11

lgG1 Calculated MW:

49 kDa Observed MW: 50-55 kDa

Purification Method:

Protein G purification CloneNo.:

2D11E5

Recommended Dilutions: WB 1:1000-1:9800 IHC 1:250-1:1000

Applications

Tested Applications: IHC, WB, ELISA

Species Specificity: Human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, Calyculin A treated HSC-T6 cells,

NIH/3T3 cells, PC-3 cells, HeLa cells

IHC: mouse testis tissue,

Background Information

STK11(serine/threonine-protein kinase 11) is also named as LKB1, PJS, and belongs to the protein kinase superfamily. It controls the activity of AMP-activated protein kinase (AMPK) family members, thereby playing a role in various processes such as cell metabolism, cell polarity, apoptosis and DNA damage response. The tumour suppressor protein LKB1 is a serine/threonine kinase that has been causally linked to Peutz-Jeghers syndrome (PJS). Defects in STK11 are a cause of Peutz-Jeghers syndrome (PJS) and defects in STK11 have been associated with testicular germ cell tumor (TGCT) and some sporadic cancers, especially lung cancers. STK11 has 2 isoforms with MW of 49 kDa and 45 kDa, and can be detected as 50-54 kDa after posttranslational modification.

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HEK-293 cells were subjected to SDS PAGE followed by western blot with 68016-1-1g (STK11/LKB1 antibody) at dilution of 1:4900 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 68016-1-Ig (STK11/LKB1 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 68016-1-1g (STK11/LKB1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse colon tissue using STK11/LKB1 antibody (68016-1-Ig, Clone: 2D11E5) at dilution of 1:400 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).