For Research Use Only

CD35 Monoclonal antibody

Catalog Number: 68033-1-Ig



Purification Method:

IHC 1:550-1:2200

IF 1:200-1:800

Basic Information

Applications

Catalog Number: GenBank Accession Number:

68033-1-lg Protein G purification NM 000573 GeneID (NCBI): CloneNo.: Size:

150ul, Concentration: 1000 µg/ml by 1378 1B8F6 Nanodrop and 549 µg/ml by Bradford Full Name: Recommended Dilutions:

method using BSA as the standard; complement component (3b/4b) receptor 1 (Knops blood group)

Mouse Calculated MW: 224 kDa Isotype:

lgG1

Immunogen Catalog Number:

Tested Applications:

AG30268

Positive Controls:

IF, IHC, ELISA IHC: human tonsillitis tissue, Species Specificity: IF: human appendicitis tissue, Human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

CD35, also known as complement receptor type 1 (CR1) or C3b/C4b receptor, is a single-chain glycoprotein consisting of 30 repeating homologous protein domains known as short consensus repeats (~60 amino acids each) followed by transmembrane and cytoplasmic domains (PMID: 8765013). CD35 is variably expressed by granulocytes, monocytes, B cells, some T cells, erythrocytes, follicular dendritic cells, Langerhans cells, and glomerular podocytes (PMID: 19004497). CD35 acts as a receptor for C3b and C4b and plays important roles in the regulation of the complement cascade and clearance of immune complexes. A soluble form of CD35 has been found in plasma (PMID: 3156931). It has been reported that soluble CD35 is produced by a proteolytic cleavage in the Cterminal region of CD35 transmembrane domain (PMID: 9405292).

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 68033-1-Ig (CD35 antibody) at dilution of 1:1100 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 68033-1-1g (CD35 antibody) at dilution of 1:1100 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immun of luorescent analysis of (4% PFA) fixed human appendicitis tissue using CD35 antibody (68033-1-lg, Clone: 188E6) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immun of luorescent analysis of (4% PFA) fixed human appendicitis tissue using CR1 antibody (68033-1-lg, Clone: 1B8E6) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).