

For Research Use Only

chemical compound m6A Monoclonal antibody



Catalog Number: 68055-1-Ig

24 Publications

Basic Information

Catalog Number:

68055-1-Ig

GenBank Accession Number:

m6A

Purification Method:

Protein A purification

Size:

150ul, Concentration: 1000 µg/ml by Nanodrop;

GeneID (NCBI):

Full Name:

CloneNo.:

1D5E10

Source:

Mouse

Recommended Dilutions:

RIP 1:1000-1:4000

IHC 1:2000-1:8000

ELISA 1:2000-1:20000

Dot Blot 1:1000-1:4000

Isotype:

IgG3

Applications

Tested Applications:

RIP, IHC, ELISA, Dot Blot

Positive Controls:

RIP: RNA,

Cited Applications:

WB, IP, IF, RIP, IHC

IHC: mouse testis tissue, human lung cancer tissue, human breast cancer tissue, human colon cancer tissue, rat lymph node

Species Specificity:

m6A, Human, Mouse, Rat

ELISA: m6A,

Cited Species:

human, rat, mouse, pig

Dot Blot: RNA,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

m6A (N6-methyladenosine) is the most abundant internal modification in mammalian mRNA. This modification is installed by the m6A methyltransferases or termed "writers" such as METTL3 and METTL14, and can be reversed by demethylases that serve as "erasers" such as FTO and ALKBH5. The stability of m6A-modified mRNA is regulated by m6A reader protein YTHDFs, which recognizes m6A and reduces the stability of target transcripts. m6A modification and its regulatory proteins play critical roles in cancer pathogenesis and progression. m6A modification is also involved in viruses life cycles, suggesting that drugs targets to m6A pathway could be used for antiviral therapy.

Protocol for Dot Blot:

<https://www.ptglab.com/protocol/68055-1-IgDotBlot.pdf>

Notable Publications

Author	Pubmed ID	Journal	Application
Lee J Martin	36359844	Cells	IHC
Meige Sun	36358640	Cancers (Basel)	
Shu Fang	35571033	Front Genet	

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Total RNA was isolated from HEK-293 cell line and was dotted to NC membrane at different amount as indicated above the dots. The membrane was blocked with BSA and blotted with m6A antibody 68055-1-Ig at 1:2000 followed by incubation of HRP-goat anti-mouse secondary antibody. Signal was developed by ECL substrate. A parallel dot blot was performed using unrelated antibody with the same isotype (UCP2 antibody 66700-1-Ig) at the same dose.




Indirect ELISA and competitive ELISA results show that this antibody is specific to m6A. Indirect ELISA was performed by coating BSA conjugated m6A at 20ng/well followed by blocking with 1% BSA. Serial diluted primary antibody was added to the plates and incubated at 37°C. HRP-goat anti-mouse was used for detection. Competitive ELISA was performed similarly except that different concentration of m6A or its structure analogue compounds are mixed in primary antibody



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 68055-1-Ig (m6A antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



HEK-293 cells were lysed and immunoprecipitated with Protein A-m6A antibody and Protein A-mouse IgG3 control antibody respectively in the presence of RNAase inhibitor cocktail. The immunoprecipitated complex was washed diggested by RNAase A followed by western blot with YTHDF1(m6A reader) antibody 17479-1-AP (1:2000). (Lysate: 3.6mg per IP; IP: 15µg antibody and 50µL beads, 4 hours at 4°C; Diggestion: 50µg/mL * 80µL RNAse A for 1



HEK-293 cells were lysed and immunoprecipitated with Protein A-m6A antibody and Protein A-mouse IgG3 control antibody respectively in the presence of RNAase inhibitor cocktail. The immunoprecipitated complex was washed diggested by RNAase A followed by western blot with IGF2BP3 (m6A reader) antibody 66526-1-Ig (1:2000). (Lysate: 4.0 mg per IP; IP: 30µg antibody and 50µL beads, 4 hours at 4°C; Diggestion: 50µg/mL * 80µL RNAse A for 1