For Research Use Only

Septin 5 Monoclonal antibody

Catalog Number: 68075-1-Ig



Basic Information

Catalog Number: GenBank Accession Number:

68075-1-lg BC025261
Size: GeneID (NCBI):

150ul , Concentration: 1000 ug/ml by 5413

Nanodrop; UNIPROT ID: Source: Q99719

Mouse Full Name:

Isotype:septin 5IgG1Calculated MW:Immunogen Catalog Number:369 aa, 43 kDaAG2205Observed MW:

40-43 kDa

Purification Method:

Protein G purification

CloneNo.: 1E2A3

Recommended Dilutions:

WB: 1:5000-1:50000

FC (Intra): 0.25 ug per 10^6 cells in a

100 μl suspension

Applications

Tested Applications: WB, FC (Intra), ELISA

Species Specificity:

human, mouse, rat, pig, rabbit

Positive Controls:

WB: pig brain tissue, LNCaP cells, Jurkat cells, K-562 cells, rabbit brain tissue, rat brain tissue, mouse brain

tissue

FC (Intra): Jurkat cells,

Background Information

Septin 5, a septin family member formerly called cell division control related protein 1 (CDCrel), is encoded in the 200 kb region and is abundantly expressed in the developing and adult brains of mice and humans. Although its constitutive deletion does not cause detectible structural or developmental abnormalities in the mouse brain, Septin 5 has been implicated in subtle neuronal and behavioral functions. Septin 5 is expressed in presynaptic terminals and exerts negative regulatory control over neurotransmitter release by binding to SNARE-protein, which has been implicated in neuropsychiatric disorders (PMID: 19240081). Septin 5 has 2 isoforms with the molecular mass of 39 and 43 kDa.

Storage

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68075-1-lg (Septin 5 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



1x10^6 Jurkat cells were intracellularly stained with 0.25 ug Septin 5 Monoclonal antibody (68075-1-lg, Clone:1E2A3) and Coralite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1)(red), or 0.25 ug Mouse IgG1 isotype control Mouse McAb (66360-1-lg, Clone: 1F8D3) (blue). Cells were fixed with 49% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).