For Research Use Only

BAG3 Monoclonal antibody

Catalog Number:68076-1-lg 2 Publications



Basic Information

Catalog Number: GenBank Accession Number:

68076-1-lg BC006418
Size: Genel D (NCBI):

150ul , Concentration: 1000 ug/ml by 9531
Nanodrop; UNIPROT ID:
Source: 095817
Mouse Full Name:

Isotype: BCL2-associated athanogene 3

IgG1 Calculated MW:

Immunogen Catalog Number:61 kDaAG17608Observed MW:

74 kDa

Applications

Tested Applications: WB, IF/ICC, ELISA

Cited Applications: WB, IHC, IF

Species Specificity:

human, mouse, rat, pig, rabbit

Cited Species: human, rat

Positive Controls:

WB: NIH/3T3 cells, LNCaP cells, rabbit heart tissue,

Purification Method:

CloneNo.:

4G6C2

Protein A purification

Recommended Dilutions:

WB 1:5000-1:50000

IF/ICC 1:500-1:2000

HeLa cells, pig heart, rat heart

IF/ICC: HepG2 cells,

Background Information

BAG3 (Bcl2-associated athanogene 3) belongs to the BAG protein family, the co-chaperone that binds to Hsc70/Hsp70 through the BAG domain and modulates their activity in polypeptide folding. BAG3 contains also a WW domain and a proline-rich (PXXP) repeat, that mediate binding to partners different from Hsp70. Through interacting with different molecular partner, BAG3 influences several cell processes, such as apoptosis, autophagy and cell motility. BAG3 protein has been reported to sustain cell survival, resistance to therapy, and/or motility and metastatization in several tumor types, thus being identified as a potential target for anticancer therapies. In addition, defects in BAG3 are the cause of some myopathy.

Notable Publications

Author	Pubmed ID	Journal	Application
Xin Lu	35253089	J Physiol Biochem	IHC
Jiao Lu	37990150	J Physiol Sci	WB,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68076-1-1g (BAG3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using BAG3 antibody (68076-1-Ig, Clone: 4G6C2) at dilution of 1:1000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).