## For Research Use Only

## C1QBP Monoclonal antibody

Catalog Number:68084-1-lg Featured Product



**Basic Information** 

Catalog Number: GenBank Accession Number:

68084-1-lg BC013731 GeneID (NCBI): Size:

150ul, Concentration: 1000 ug/ml by 708 Nanodrop: **UNIPROT ID:** Q07021 Mouse Full Name:

Isotype: complement component 1, q lgG2b subcomponent binding protein

Immunogen Catalog Number: Calculated MW: 282 aa, 31 kDa AG19773

> Observed MW: 32 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, FC (Intra), ELISA

Species Specificity: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: LNCaP cells, HEK-293 cells, mouse brain tissue, HeLa cells, HepG2 cells, Jurkat cells, K-562 cells

**Purification Method:** 

CloneNo.:

1F9B1

Protein A purification

Recommended Dilutions:

WB 1:5000-1:50000

IF/ICC 1:400-1:1600

IHC 1:500-1:2000

IHC: human tonsillitis tissue, human liver cancer

IF/ICC: HeLa cells,

## **Background Information**

initially copurified with splicing factor SF2 (PMID: 1830244). The protein is synthesized as a pro-protein of 282 amino acids (aa) that is post-translationally processed by removal of the initial 73 aa to a mature protein of 209 aa (PMID: 8262387). C1QBP is an evolutionary conserved and ubiquitously expressed multifunctional protein and has been reported to be a predominantly mitochondrial matrix protein involved in inflammation and infection  $processes, mitochondrial\ ribosome\ biogenesis, regulation\ of\ apoptosis\ and\ nuclear\ transcription,\ and\ pre-mRNA$ splicing (PMID: 28942965).

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com


## Selected Validation Data




WB result of C1QBP antibody (68084-1-Ig; 1:6000; incubated at room temperature for 1.5 hours) with sh-Control and sh-C1QBP transfected HEK-293 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 68084-1-Ig (C1QBP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.




Immunofluorescent analysis of (-20°C Methanol) fixed Hela cells using C1QBP antibody (68084-1-lg, Clone: 1F9B1) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 68084-1-1g (C1QBP antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 68084-1-lg (C1QBP antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10^6 HeLa cells were intracellularly stained with 0.5 ug Anti-Human C1QBP (68084-1-lg, Clone:1F9B1) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgC(H+L) at dilution 1:1000 (red), or 0.5 ug Mouse IgC2b Isotype Control (66360-3-lg, Clone: K11B8C4B5) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).