## For Research Use Only

## Annexin VI Monoclonal antibody

Catalog Number: 68086-1-Ig



**Basic Information** 

Catalog Number: GenBank Accession Number:

68086-1-lg BC017046 GeneID (NCBI):

150ul, Concentration: 500 ug/ml by

Nanodrop: **UNIPROT ID:** 

P08133 Mouse Full Name:

Isotype: annexin A6 lgG1 Calculated MW:

Immunogen Catalog Number: 673 aa, 76 kDa AG3263 Observed MW:

67 kDa

**Purification Method:** 

Protein G purification

CloneNo.: 2H7G7

Recommended Dilutions:

WB: 1:2000-1:10000 IF/ICC: 1:200-1:800

FC (Intra): 0.40 ug per 10^6 cells in a

100 µl suspension

**Applications** 

**Tested Applications:** 

WB, IF/ICC, FC (Intra), ELISA

Species Specificity:

human, mouse, rat, pig, rabbit

**Positive Controls:** 

WB: LNCaP cells, HeLa cells, human placenta tissue, pig brain tissue, rabbit brain tissue, rat brain tissue,

mouse brain tissue IF/ICC: HepG2 cells,

FC (Intra): K-562 cells,

## **Background Information**

ANXA6, also called CBP68, or 67 kDa calelectrin, belongs to a family of calcium-dependent membrane and phospholipid binding proteins. Several members of the annexin family have been implicated in membrane-related events along exocytotic and endocytotic pathways. The annexin VI gene is approximately 60 kbp long and contains 26 exons. It encodes a protein of about 68 kDa that consists of eight 68-amino acid repeats separated by linking  $sequences\ of\ variable\ lengths.\ It\ is\ highly\ similar\ to\ human\ annexins\ l\ and\ ll\ sequences,\ each\ of\ which\ contain\ four$ such repeats. Annexin VI has been implicated in mediating the endosome aggregation and vesicle fusion in secreting epithelia during exocytosis.

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 68086-1-lg (Annexin VI antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using Annexin VI antibody (68086-1-Ig, Clone: 2H7G7) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L).



1X10^6 K-562 cells were intracellularly stained with 0.4 ug Anti-Human Annexin VI (68086-1-1g, Clone:2H7G7) and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(I+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-1g, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).