For Research Use Only

Annexin VI Monoclonal antibody

Catalog Number: 68086-1-Ig



Basic Information

Catalog Number: GenBank Accession Number: BC017046

68086-1-lg GeneID (NCBI):

150ul, Concentration: 500 ug/ml by

Nanodrop: **UNIPROT ID:**

P08133 Mouse Full Name: Isotype: annexin A6 lgG1 Calculated MW: Immunogen Catalog Number: 673 aa, 76 kDa

AG3263 Observed MW:

67 kDa

Applications

Tested Applications:

WB, IF/ICC, FC (Intra), ELISA

Species Specificity:

human, mouse, rat, pig, rabbit

Purification Method:

Protein G purification

CloneNo.: 2H7G7

Recommended Dilutions:

WB 1:2000-1:10000 IF/ICC 1:200-1:800

WB: LNCaP cells, HeLa cells, human placenta tissue, pig brain tissue, rabbit brain tissue, rat brain tissue,

Background Information

ANXA6, also called CBP68, or 67 kDa calelectrin, belongs to a family of calcium-dependent membrane and phospholipid binding proteins. Several members of the annexin family have been implicated in membrane-related events along exocytotic and endocytotic pathways. The annexin VI gene is approximately 60 kbp long and contains 26 exons. It encodes a protein of about 68 kDa that consists of eight 68-amino acid repeats separated by linking sequences of variable lengths. It is highly similar to human annexins I and II sequences, each of which contain four such repeats. Annexin VI has been implicated in mediating the endosome aggregation and vesicle fusion in secreting epithelia during exocytosis.

Positive Controls:

mouse brain tissue IF/ICC: HepG2 cells,

Storage

Store at -20°C. Stable for one year after shipment.

Aliquoting is unnecessary for -20°C storage


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


*** 20ul sizes contain 0.1% BSA

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68086-1-lg (Annexin VI antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using Annexin VI antibody (68086-1-Ig, Clone: 2H7G7) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L).



1X10^6 K-562 cells were intracellularly stained with 0.4 ug Anti-Human Annexin VI (68086-1-1g, Clone:2H7G7) and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(I+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-1g, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).