

For Research Use Only

SMG6 Monoclonal antibody

Catalog Number: 68109-1-Ig



Basic Information

Catalog Number: 68109-1-Ig	GenBank Accession Number: NM_017575	Purification Method: Protein A purification
Size: 150ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 23293	CloneNo.: 1D2G2
Source: Mouse	Full Name: Smg-6 homolog, nonsense mediated mRNA decay factor (C. elegans)	Recommended Dilutions: WB 1:1000-1:6000
Isotype: IgG2a	Calculated MW: 160 kDa	
Immunogen Catalog Number: AG30182	Observed MW: 160 kDa	

Applications

Tested Applications:
WB, ELISA

Species Specificity:
Human, mouse

Positive Controls:

WB : LNCaP cells, RAW 264.7 cells, MCF-7 cells, HeLa cells, HEK-293 cells

Background Information

SMG6, also known as C17orf31, EST1A, and hEST1A, is a component of the telomerase ribonucleoprotein (RNP) complex that is essential for the replication of chromosome termini (PMID: 19179534). It promotes in vitro the ability of TERT to elongate telomeres (PMID: 12676087, 12699629). SMG6 also plays a role in the nonsense-mediated mRNA decay (NMD) pathway, providing the endonuclease activity near the premature translation termination codon that is needed to initiate NMD (PMID: 19060897, 20930030). Alternate splicing results in multiple transcript variants (PMID: 12699629, 14636577).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.

Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68109-1-Ig (SMG6 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with Lamin B1 Monoclonal antibody (66095-1-Ig) as loading control.



RAW 264.7 cells were subjected to SDS PAGE followed by western blot with 68109-1-Ig (SMG6 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.