For Research Use Only

LPIN1 Monoclonal antibody

Catalog Number:68124-1-lg 2 Publications



Basic Information

Catalog Number: GenBank Accession Number:

68124-1-Ig BC030537
Size: GeneID (NCBI):

150ul , Concentration: 1000 ug/ml by 23175 Nanodrop and 628 ug/ml by Bradford UNIPROT ID: method using BSA as the standard; Q14693

Source: Full Name:
Mouse lipin 1
Isotype: Calminated

Isotype: Calculated MW:
IgG2a 890 aa, 99 kDa
Immunogen Catalog Number: Observed MW:
AG25813 130 kDa

Purification Method:

Protein A purification

CloneNo.: 3G11B10

Recommended Dilutions: WB 1:2000-1:10000 IHC 1:2000-1:8000

IF/ICC 1:400-1:1600

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications:

WB, IHC

Species Specificity: Human, Rat, Mouse Cited Species:

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HepG2 cells, Jurkat cells, MOLT-4 cells, K-562

cells

IHC: human liver cancer tissue, rat skeletal muscle

tissue

IF/ICC: NIH/3T3 cells,

Background Information

LPIN1, also named as KIAA0188, belongs to the lipin family. It is a magnesium-dependent phosphatidate phosphatase enzyme which catalyzes the conversion of phosphatidic acid to diacylglycerol during triglyceride, phosphatidylcholine and phosphatidylethanolamine biosynthesis and therefore controls the metabolism of fatty acids at different levels. LPIN1 has a molecular weight of about 99 kDa but predicted bands around 130 kDa. (PMID:20231281, PMID:29765047).

Notable Publications

Author	Pubmed ID	Journal	Application
Sixin Chen	38570025	Food Chem Toxicol	WB,IHC
Shuxuan Li	38141939	Eur J Pharmacol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68124-1-1g (LPIN1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with Lamin B1 Monoclonal antibody (66095-1-1g) as loading control.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 68124-1-Ig (LPIN1 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed NIH/3T3 cells using LPIN1 antibody (68124-1-Ig, Clone: 3G11B10) at dilution of 1:800 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L).