

For Research Use Only

ACOT7 Monoclonal antibody

Catalog Number: 68140-1-Ig



Basic Information

Catalog Number: 68140-1-Ig	GenBank Accession Number: BC017365	Purification Method: Protein A purification
Size: 150ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 11332	CloneNo.: 1E9C7
Source: Mouse	UNIPROT ID: O00154	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG2a	Full Name: acyl-CoA thioesterase 7	
Immunogen Catalog Number: AG8975	Calculated MW: 370 aa, 40 kDa	
	Observed MW: 37 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : LNCaP cells, HeLa cells, HEK-293 cells, Jurkat cells, K-562 cells, HSC-T6 cells, NIH/3T3 cells, pig brain tissue, rabbit brain tissue
Species Specificity: Human, Rat, Mouse, Rabbit, Pig	

Background Information

Acyl-CoA thioesterases (ACOTs) are enzymes that catalyze hydrolysis of fatty acyl-CoAs to free fatty acids and CoA-SH. ACOTs are expressed ubiquitously in prokaryotes and eukaryotes (PMID: 11755680). ACOT7 also known as BACH, CTE-II, ACT, ACH1, BACHa, is one of the most extensively studied ACOTs as a major isoform of the ACOT family (PMID: 17563367). The enzyme is localized to the cytosol and is highly expressed in brain tissue and testis tissue, showing a preference for long-chain acyl-CoA substrates with fatty acid chains of 8-16 carbon atoms (C8-C16) (PMID: 28518146). It has been identified that ACOT7 plays a role in neurotoxic prevention and is a candidate drug target in inflammatory disease (PMID: 17563367, 23459938, 10509865). ACOT7 has 7 isoforms as molecular between 27-40 kDa, it may also form dimer (~70 kDa) (PMID: 20470824).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68140-1-Ig (ACOT7 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.