

For Research Use Only

WNK1 Monoclonal antibody

Catalog Number: 68149-1-Ig



Basic Information

Catalog Number: 68149-1-Ig	GenBank Accession Number: BC021121	Purification Method: Protein A purification
Size: 150ul , Concentration: 1000 µg/ml by Nanodrop and 638 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 65125	CloneNo.: 1B6A10
Source: Mouse	Full Name: WNK lysine deficient protein kinase 1	Recommended Dilutions: WB 1:1000-1:5000
Isotype: IgG2a	Calculated MW: 251 kDa	
Immunogen Catalog Number: AG28044	Observed MW: 250 kDa	

Applications

Tested Applications: FC, WB, ELISA	Positive Controls: WB : HeLa cells, HEK-293 cells, HepG2 cells, A549 cells, MCF-7 cells, Jurkat cells, HSC-T6 cells, NIH/3T3 cells
Species Specificity: Human, Mouse, Rat	

Background Information

WNK kinase is a unique serine-threonine kinase in which the lysine (K) residue in the catalytic domain, that is found in the majority of kinases is not conserved. In mammals, there are four WNK kinase family members (WNK1-4). WNK1 has 6 isoforms, such as long form WNK1 (L-WNK1, 250 kDa, strong expression in dorsal root ganglia and spinal cord); kidney-specific isoform (KS-WNK1, 207 kDa, lacks a kinase domain). KS-WNK1 cannot phosphorylate substrates and instead interacts with and modulates the actions of other WNKs via a carboxyl terminal HQ domain.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68149-1-Ig (WNK1 antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.



1×10^6 HEK-293T cells were intracellularly stained with 0.4 μ g Anti-Human WNK1 (68149-1-Ig, Clone:1B6A10) and CoraLite[®]488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 μ g Mouse IgG2a Isotype Control (C1.18.4) (65208-1-Ig, Clone: C1.18.4) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).