For Research Use Only

DDAH2 Monoclonal antibody

Catalog Number: 68154-1-Ig



Basic Information

Catalog Number: GenBank Accession Number:

68154-1-lg BC001435
Size: GeneID (NCBI):

150ul , Concentration: 1000 ug/ml by 23564
Nanodrop; UNIPROT ID:
Source: 095865

Mouse Full Name:
Isotype: dimethylarginine

IgG3 dimethylaminohydrolase 2

Immunogen Catalog Number:Calculated MW:AG712729 kDa

Observed MW: 30 kDa

Applications

Tested Applications:

WB, ELISA
Species Specificity:
Human, Rat, Mouse, Rabbit

Positive Controls:

WB: rabbit brain tissue, mouse retina tissue, rat colon tissue, rat brain tissue, mouse brain tissue

Purification Method:

CloneNo.:

1A10G3

Protein A purification

Recommended Dilutions:

WB 1:5000-1:50000

Background Information

The DDAH family of proteins is formed by two isoforms, DDAH-1 and DDAH-2, both of which utilize ADMA as their substrate and increase NOS activity. DDAH-2, also named as DDAH, G6A, NG30, is the most prevalent isoform in blood vessels and endothelium and is important for embryonic development. It has also been linked to other diseases, such as hypertension, cerebral hemorrhage, cardiac disease, and diabetes. In cultured human endothelial cells, DDAH-2 had been found only in the cytosol, but in the nuclei of rat vascular smooth muscle cells.

Storage

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68154-1-1g (DDAH2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



mouse retina tissue were subjected to SDS PAGE followed by western blot with 68154-1-1g (DDAH2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



rat colon tissue were subjected to SDS PAGE followed by western blot with 68154-1-1g (DDAH2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.