

For Research Use Only

# U2AF2/U2AF65 Monoclonal antibody, PBS Only



Catalog Number: 68166-1-PBS

## Basic Information

<b>Catalog Number:</b> 68166-1-PBS	<b>GenBank Accession Number:</b> BC008740	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 100ug , Concentration: 1mg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 11338	<b>CloneNo.:</b> 1B7E7
<b>Source:</b> Mouse	<b>UNIPROT ID:</b> P26368	
<b>Isotype:</b> IgG2a	<b>Full Name:</b> U2 small nuclear RNA auxiliary factor 2	
<b>Immunogen Catalog Number:</b> AG8070	<b>Calculated MW:</b> 54 kDa	
	<b>Observed MW:</b> 60-65 kDa	

## Applications

**Tested Applications:**  
WB, IHC, IF/ICC, Indirect ELISA

**Species Specificity:**  
human, mouse, rat

## Background Information

U2AF2, also named as U2 auxiliary factor 65 kDa subunit, is a 475 amino acid protein, which belongs to the splicing factor SR family. U2AF2 is Necessary for the splicing of pre-mRNA. By recruiting PRPF19 and the PRP19C/Prp19 complex/NTC/Nineteen complex to the RNA polymerase II C-terminal domain (CTD), pre-mRNA may couple transcription to splicing. The calculated molecular weight of U2AF2 is 53 kDa, but modified U2AF2 is about 65 kDa.

## Storage


**Storage:**  
Store at -80°C.

**Storage Buffer:**  
PBS Only


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 68166-1-Ig (U2AF2 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 68166-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using U2AF2 antibody (68166-1-Ig, Clone: 1B7E7) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 68166-1-PBS in a different storage buffer formulation.




1X10<sup>6</sup> HEK-293 cells were intracellularly stained with 0.4 ug Anti-Human U2AF2 (68166-1-Ig, Clone:1B7E7) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2a Isotype Control (C1.18.4) (65208-1-Ig, Clone: C1.18.4) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 68166-1-PBS in a different storage buffer




Various lysates were subjected to SDS PAGE followed by western blot with 68166-1-Ig (U2AF2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control. This data was developed using the same antibody clone with 68166-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 68166-1-Ig (U2AF2 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 68166-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded rat liver tissue slide using 68166-1-Ig (U2AF2 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 68166-1-PBS in a different storage buffer formulation.



4T1 cells were subjected to SDS PAGE followed by western blot with 68166-1-Ig (U2AF2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68166-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using U2AF2 antibody (68166-1-Ig, Clone: 1B7E7) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 68166-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using U2AF2 antibody (68166-1-Ig, Clone: 1B7E7) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 68166-1-PBS in a different storage buffer formulation.