## For Research Use Only

# **TOLLIP Monoclonal antibody**

Catalog Number:68170-1-lg Featured Product



**Basic Information** 

Catalog Number: GenBank Accession Number:

68170-1-lg BC012057 GeneID (NCBI): Size: 150ul , Concentration: 1000 ug/ml by 54472

Nanodrop: **UNIPROT ID:** Q9H0E2 Mouse Full Name:

Isotype: toll interacting protein

lgG1 Calculated MW: Immunogen Catalog Number: 30 kDa AG32731

Observed MW:

30 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, FC (Intra), ELISA

Species Specificity: human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### **Positive Controls:**

WB: pig brain tissue, LNCaP cells, HepG2 cells, Jurkat cells, K-562 cells, human peripheral blood platelets, rabbit brain tissue, rat brain tissue, mouse brain tissue

**Purification Method:** 

CloneNo.:

1B3C4

Protein G purification

Recommended Dilutions:

WB 1:5000-1:50000

IF/ICC 1:400-1:1600

IHC 1:400-1:1600

IHC: mouse colon tissue, rat colon tissue

IF/ICC: U2OS cells. FC (Intra): Jurkat cells,

# **Background Information**

 $TOLLIP\ (Toll\ interacting\ protein), is\ an\ inhibitory\ adaptor\ protein.\ Tollip\ is\ a\ negative\ regulator\ of\ IL-1R\ and\ Toll-like$  $receptors\ signal ling\ pathway, involved\ in\ inflammatory\ immune\ response\ and\ inhibiting\ IRAK1\ phosphory lation$ and kinase activity (PMID:11441107, PMID:10854325, PMID:11751856). The TOLLIP-dependent selective autophagy pathway plays a role in clearance of cytotoxic polyQ proteins aggregates (PubMed:25042851).

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


### Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68170-1-lg (TOLLIP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 68170-1-1g (TOLLIP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control




WB result of TOLLIP antibody (68170-1-lg; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-TOLLIP transfected HepG2 cells.




Immunohistochemical analysis of paraffinembedded rat colon tissue slide using 68170-1-Ig (TOLLIP antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse colon tissue slide using 68170-1-1g (TOLLIP antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse colon tissue slide using 68170-1-1g (TOLLIP antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed U2OS cells using TOLLIP antibody (68170-1-lg, Clone: 183C4) at dilution of 1:200 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse  $\lg G(H+L)$ .



1X10^6 Jurkat cells were intracellularly stained with 0.4 ug Anti-Human TOLLIP (68170-1-1g, Clone:1B3C4) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-1g, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).