

For Research Use Only



APPL1 Monoclonal antibody, PBS Only

Catalog Number: 68195-1-PBS

Featured Product

Basic Information

Catalog Number:

68195-1-PBS

Size:

100ug, Concentration: 1mg/ml by Nanodrop;

Source:

Mouse

Isotype:

IgG2b

Immunogen Catalog Number:

AG3334

GenBank Accession Number:

BC028599

GeneID (NCBI):

26060

UNIPROT ID:

Q9UKG1

Full Name:

adaptor protein, phosphotyrosine interaction, PH domain and leucine zipper containing 1

Calculated MW:

709 aa, 80 kDa

Observed MW:

80 kDa

Purification Method:

Protein A purification

CloneNo.:

1B7B11

Applications

Tested Applications:

Indirect ELISA, IF/ICC, IHC, WB

Species Specificity:

rabbit, pig, rat, mouse, human

Background Information

Adaptor protein, phosphotyrosine interaction, PH domain and leucine zipper containing 1 (APPL1), a binding partner of Akt2 and an important regulator of INS signaling, plays a key role in the regulation of INS secretion [PMID:22615370]. APPL1 interacts with adiponectin receptors and mediates the INS-sensitizing effects of adiponectin in muscle and endothelial cells. It also participates in nuclear signaling and transcriptional regulation, mostly by modulating the activity of various nuclear factors [PMID:22685329]. Apart from its role in endocytosis and endosomal transport, APPL1 was reported to undergo nucleocytoplasmic shuttling and participate in transcriptional regulation, e.g. by interactions with histone deacetylases (HDACs) [PMID:19686092].

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS Only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68195-1-Ig (APPL1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68195-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 68195-1-Ig (APPL1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68195-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 68195-1-Ig (APPL1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68195-1-PBS in a different storage buffer formulation.




HepG2 cells were subjected to SDS PAGE followed by western blot with 68195-1-Ig (APPL1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68195-1-PBS in a different storage buffer formulation.




pig heart tissue were subjected to SDS PAGE followed by western blot with 68195-1-Ig (APPL1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68195-1-PBS in a different storage buffer formulation.




WB result of APPL1 antibody (68195-1-Ig; 1:30000; incubated at room temperature for 1.5 hours) with sh-Control and sh-APPL1 transfected HEK-293 cells. This data was developed using the same antibody clone with 68195-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 68195-1-Ig (APPL1 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 68195-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using APPL1 antibody (68195-1-Ig, Clone: 1B7B11) at dilution of 1:400 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 68195-1-PBS in a different storage buffer formulation.



1X10⁶ HepG2 cells were intracellularly stained with 0.4 ug Anti-Human APPL1 (68195-1-Ig, Clone:1B7B11) and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2b Isotype Control (MPC-11) (65128-1-Ig, Clone: MPC-11) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 68195-1-PBS in a