

For Research Use Only

CYP2S1 Monoclonal antibody

Catalog Number: 68198-1-Ig



Basic Information

Catalog Number: 68198-1-Ig	GenBank Accession Number: BC033691	Purification Method: Protein G purification
Size: 150ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 29785	CloneNo.: 3E5D4
Source: Mouse	UNIPROT ID: Q96SQ9	Recommended Dilutions: WB 1:2000-1:10000
Isotype: IgG1	Full Name: cytochrome P450, family 2, subfamily S, polypeptide 1	
Immunogen Catalog Number: AG4605	Calculated MW: 504 aa, 56 kDa	
	Observed MW: 50-56 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : NCCIT cells, HeLa cells, A549 cells, pig colon tissue, pig rectum tissue
Species Specificity: Human, Pig	

Background Information

The cytochrome P450 (CYP) family of isoenzymes are involved in the metabolism of diverse endogenous compounds and detoxification of exogenous agents. Cytochrome P450 2S1 (CYP2S1) is one 'orphan' CYP, and is particularly expressed in epithelial cells of tissues which are exposed to the environment such as the skin, respiratory, gastrointestinal, and urinary tracts. CYP2S1 is associated primarily with the synthesis and metabolism of lipids including prostaglandins (PGE2) and retinoids. In addition, CYP2S1 expression is associated with patient survival in breast cancer.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68198-1-Ig (CYP2S1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



Various lysates were subjected to SDS PAGE followed by western blot with 68198-1-Ig (CYP2S1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.