

For Research Use Only

ARF4 Monoclonal antibody

Catalog Number: 68204-1-Ig **Featured Product**



Basic Information

Catalog Number: 68204-1-Ig	GenBank Accession Number: BC003364	Purification Method: Protein A purification
Size: 150ul , Concentration: 1000 µg/ml by Nanodrop and 640 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 378	CloneNo.: 2A4B8
Source: Mouse	Full Name: ADP-ribosylation factor 4	Recommended Dilutions: WB 1:5000-1:50000 IF 1:400-1:1600
Isotype: IgG2b	Calculated MW: 180 aa, 21 kDa	
Immunogen Catalog Number: AG25522	Observed MW: 18 kDa	

Applications

Tested Applications: FC, IF, WB, ELISA	Positive Controls:
Species Specificity: Human, mouse, rat	WB : LNCaP cells, HEK-293 cells, rat brain tissue, A549 cells, mouse brain tissue, HeLa cells, HepG2 cells, Jurkat cells
	IF : HepG2 cells,

Background Information

ARF4, also named as ARF2, belongs to the small GTPase superfamily and Arf family. ADP-ribosylation factors (ARFs) are members of the ARF family of GTP-binding proteins of the Ras superfamily, with 20 kDa protein size. ARFs bind and regulate GTP/GDP cycle by alternating between the active GTP-bound and inactive GDP-bound conformations. ARF family proteins are essential and ubiquitous in eukaryotes. Six highly conserved members of the family have been identified in mammalian cells. They function in vesicular traffic and actin remodelling and other bioprocesses in cells. ARF4 and ARF5 are class II ARFs. Their function are not fully described. (PMID: 7759471, PMID: 16042562). This antibody can bind ARFs for the close sequences.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68204-1-Ig (ARF4 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




A549 cells were subjected to SDS PAGE followed by western blot with 68204-1-Ig (ARF4 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




WB result of ARF4 antibody (68204-1-Ig; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ARF4 transfected HeLa cells.




Various lysates were subjected to SDS PAGE followed by western blot with 68204-1-Ig (ARF4 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 68204-1-Ig (ARF4 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using ARF4 antibody (68204-1-Ig, Clone: 2A4B8) at dilution of 1:800 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



1×10^6 K-562 cells were intracellularly stained with 0.4 ug Anti-Human ARF4 (68204-1-Ig, Clone:2A4B8) and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2b Isotype Control (66360-3-Ig, Clone: K11B8C4B5) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).