For Research Use Only

DNM2 Monoclonal antibody

Catalog Number: 68209-1-Ig



Basic Information

Catalog Number: GenBank Accession Number:

 68209-1-Ig
 BC054501
 Protein G purification

 Size:
 GeneID (NCBI):
 CloneNo.:

 150ul . Concentration: 1000 ug/ml by 1785
 188H3

150ul , Concentration: 1000 µg/ml by 1785
Nanodrop; Full Name:
Source: dynamin 2
Mouse Calculated MW:

Isotype: 98 kDa
IgG1 Observed MW:
Immunogen Catalog Number: 100 kDa

AG6274

Applications
Tested Applications:

IF, WB, ELISA
Species Specificity:

Human, mouse, rat, pig, rabbit

Positive Controls:

WB: HCT 116 cells, HeLa cells, HEK-293 cells, Jurkat cells, K-562 cells, pig brain tissue, rabbit brain tissue, rat brain tissue, mouse brain tissue

Purification Method:

Recommended Dilutions:

WB 1:5000-1:50000

IF 1:200-1:800

IF · Hel a cells

Background Information

DNM2, also known as DYN2, DYNII, CMTDI1, CMTDIB, belongs to the dynamin family. DNM2 is a ubiquitously expressed large GTPase involved in clathrin-dependent and -independent endocytosis and intracellular membrane trafficking. DNM2 plays a role in the regulation of neuron morphology, axon growth, and the formation of neuronal growth cones. Dynamins are associated with microtubules. DNM2 plays an important role in endocytosis and is involved in cytokinesis (PMID: 33713620, 12498685). Mutations in DNM2 cause dominant centronuclear myopathy (PMID: 16227997). Defects in DNM2 are the cause of Charcot-Marie-Tooth disease dominant intermediate type B (CMTDIB) (PMID: 15731758). Alternate splicing results in multiple transcript variants.

Storage

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68209-1-lg (DNM2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.



Immunofluorescent analysis of (-20°C Methanol) fixed Hela cells using DNM2 antibody (68209-1-Ig, Clone: 1B8H3) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).