For Research Use Only

ACTN2 Monoclonal antibody

Catalog Number:68223-1-lg 1 Publications



Basic Information

Catalog Number: GenBank Accession Number:

68223-1-lg BC051770

ize: GeneID (NCBI):

150ul, Concentration: 1000 ug/ml by 88

Nanodrop; UNIPROT ID: Source: P35609

Mouse Full Name:
Isotype: actinin, alpha 2
IgG2b Calculated MW:
Immunogen Catalog Number: 104 kDa

AG5578 Observed MW:

103 kDa

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity:

Human, mouse, rat, pig, rabbit

Cited Species: mouse Positive Controls:

WB: pig heart tissue, rat colon tissue, rabbit heart tissue, rat heart tissue, mouse heart tissue, pig colon

Purification Method:

Protein A purification

Recommended Dilutions:

WB 1:20000-1:100000

CloneNo.:

1D5F9

tissu

Background Information

Alpha actinin 2 (ACTN2) belongs to the alpha-actinin family and is expressed in both skeletal and cardiac muscles and functions to anchor myofibrillar actin thin filaments and titin to Z-discs (PMID: 30701273). ACTN2 is an actin-binding protein with multiple roles in different cell types. In nonmuscle cells, the cytoskeletal isoform is found along microfilament bundles and adherens-type junctions, where it is involved in binding actin to the membrane. In contrast, skeletal, cardiac, and smooth muscle isoforms are localized to the Z disc and analogous dense bodies, where they help anchor the myofibrillar actin filaments. Mutations in ACTN2 are associated with hypertrophic cardiomyopathy, as well as dilated cardiomyopathy and endocardial fibroelastosis (PMID: 20022194, 14567970).

Notable Publications

Author	Pubmed ID	Journal	Application
Shan Gao	38495825	iScience	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68223-1-1g (ACTN2 antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.



rat colon tissue were subjected to SDS PAGE followed by western blot with 68223-1-lg (ACTN2 antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.