For Research Use Only

IMMT Monoclonal antibody

Catalog Number:68226-1-lg Featured Product



Purification Method:

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number:

68226-1-lg Protein A purification BC002412 GeneID (NCBI): CloneNo.:

150ul, Concentration: 500 µg/ml by 10989 2D2F8

Source: inner membrane protein, WB 1:5000-1:50000 mitochondrial (mitofilin) IHC 1:1000-1:4000 Mouse IF 1:400-1:1600

Calculated MW: Isotype: 90 kDa lgG2b Immunogen Catalog Number: Observed MW: 90 kDa

Applications

Tested Applications: Positive Controls:

FC, IF, IHC, WB, ELISA WB: HeLa cells, HepG2 cells Species Specificity: IHC: mouse heart tissue, Human, Rat, Mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

IMMT (also known as mitofilin) is an inner mitochondrial membrane protein that is preferentially expressed in heart tissue and is commonly used as the marker for mitochondria. Mitofilin is known to be a critical organizer of mitochondrial cristae morphology and is indispensable for normal mitochondrial function. Three isoforms of mitofilin exist due to the alternative splicing. All three proteins of 87kD, 89kD and 80kD can be detected in immunoblot analysis with this antibody.

IF: HeLa cells,

Storage

Storage:

Nanodrop;

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68226-1-lg (IMMT antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 68226-1-1g (IMMT antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 68226-1-Ig (IMMT antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed Hela cells using IMMT antibody (68226-1-lg, Clone: 2D2E8) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using IMMT antibody (68226-1-lg, Clone: 2D2E8) at dilution of 1:800 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).



1X10^6 HEK-293 cells were intracellularly stained with 0.4 ug Anti-Human IMMT (68226-1-lg, Clone:2D2E8) and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2b Isotype Control (66360-3-lg, Clone: K11B8C4B5) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



WB result of IMMT antibody (68226-1-lg; 1:30000; incubated at room temperature for 1.5 hours) with sh-Control and sh-IMMT transfected HeLa cells.