For Research Use Only

IMMT Monoclonal antibody

Catalog Number:68226-1-lg Featured Product 1 Publications


oroteintech Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 68226-1-lg	GenBank Accession Numbe BC002412	r: Purification Method: Protein A purification
	Size:	GenelD (NCBI):	CloneNo.:
	150ul , Concentration: 500 ug/ml by	10989	2D2E8
	Nanodrop;	UNIPROT ID:	Recommended Dilutions:
	Source:	Q16891	WB 1:5000-1:50000
	Mouse	Full Name:	IHC 1:1000-1:4000 IF/ICC 1:400-1:1600
	Isotype: IgG2b Immunogen Catalog Number: AG0102	inner membrane protein, mitochondrial (mitofilin)	11/100 1.400-1.1000
		Calculated MW:	
		90 kDa Observed MW:	
		90 kDa	
Applications	Tested Applications:		
	WB, IHC, IF/ICC, FC (Intra), ELISA	WB:	HeLa cells, HepG2 cells
	Species Specificity:	IHC	: mouse heart tissue,
	human, mouse, rat	IF/I	CC : HeLa cells,
	Note-IHC: suggested antic with TE buffer pH 9.0; (*) A antigen retrieval may be p with citrate buffer pH 6.0	Alternatively,	
	IMMT (also known as mitofilin) is an inner mitochondrial membrane protein that is preferentially expressed in hear tissue and is commonly used as the marker for mitochondria. Mitofilin is known to be a critical organizer of mitochondrial cristae morphology and is indispensable for normal mitochondrial function. Three isoforms of mitofilin exist due to the alternative splicing. All three proteins of 87kD, 89kD and 80kD can be detected in immunoblot analysis with this antibody.		
Background Information	tissue and is commonly used as the r mitochondrial cristae morphology ar mitofilin exist due to the alternative	narker for mitochondria. Mitc nd is indispensable for norma splicing. All three proteins o	ofilin is known to be a critical organizer of l mitochondrial function. Three isoforms of
Background Information Notable Publications	tissue and is commonly used as the r mitochondrial cristae morphology ar mitofilin exist due to the alternative immunoblot analysis with this antib	narker for mitochondria. Mitc nd is indispensable for norma splicing. All three proteins o	ofilin is known to be a critical organizer of l mitochondrial function. Three isoforms of
<u> </u>	tissue and is commonly used as the r mitochondrial cristae morphology ar mitofilin exist due to the alternative immunoblot analysis with this antib Author Put	narker for mitochondria. Mito nd is indispensable for norma splicing. All three proteins o ody.	ofilin is known to be a critical organizer of l mitochondrial function. Three isoforms of f 87kD, 89kD and 80kD can be detected in
Notable Publications	tissue and is commonly used as the r mitochondrial cristae morphology ar mitofilin exist due to the alternative immunoblot analysis with this antib Author Put Keun Woo Ryu 395 Storage: Storage Store at -20°C. Stable for one year aft Storage Buffer:	narker for mitochondria. Mito nd is indispensable for norma splicing. All three proteins o ody. omed ID Journal 506109 Nature	ofilin is known to be a critical organizer of l mitochondrial function. Three isoforms of f 87kD, 89kD and 80kD can be detected in
Notable Publications	tissue and is commonly used as the r mitochondrial cristae morphology ar mitofilin exist due to the alternative immunoblot analysis with this antib Author Put Keun Woo Ryu 399 Storage: Storage Store at -20°C. Stable for one year aff Storage Buffer: PBS with 0.02% sodium azide and 500	narker for mitochondria. Mito nd is indispensable for norma splicing. All three proteins o ody. med ID Journal 506109 Nature eer shipment.	ofilin is known to be a critical organizer of l mitochondrial function. Three isoforms of f 87kD, 89kD and 80kD can be detected in
<u> </u>	tissue and is commonly used as the r mitochondrial cristae morphology ar mitofilin exist due to the alternative immunoblot analysis with this antib Author Put Keun Woo Ryu 395 Storage: Storage Store at -20°C. Stable for one year aft Storage Buffer:	narker for mitochondria. Mito nd is indispensable for norma splicing. All three proteins o ody. med ID Journal 506109 Nature eer shipment.	ofilin is known to be a critical organizer of l mitochondrial function. Three isoforms of f 87kD, 89kD and 80kD can be detected in

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data







Various lysates were subjected to SDS PAGE followed by western blot with 68226-1-lg (IMMT antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 68226-1-Ig (IMMT antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 68226-1-Ig (IMMT antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed HeLa cells using IMMT antibody (68226-1-1g, Clone: 2D2E8) at dilution of 1:800 and CoraLite@488-Conjugated Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using IMMT antibody (68226-1-1g, Clone: 2D2E8) at dilution of 1:800 and CoraLite@488-Conjugated Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).



1X10^6 HEK-293 cells were intracellularly stained with 0.4 ug Anti-Human IMMT (68226-1-lg, Clone:2D2E8) and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2b Isotype Control (66360-3-lg, Clone: K11B8C4B5) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



WB result of IMMT antibody (68226-1-lg; 1:30000; incubated at room temperature for 1.5 hours) with sh-Control and sh-IMMT transfected HeLa cells.