## For Research Use Only

## IMMT Monoclonal antibody, PBS Only

Catalog Number: 68226-1-PBS

**Featured Product** 



**Purification Method:** 

Protein A purification

CloneNo.:

2D2E8

**Basic Information** 

Catalog Number: 68226-1-PBS

GenBank Accession Number:

BC002412

GeneID (NCBI):

100ug, Concentration: 1 mg/ml by

10989

Nanodrop:

**UNIPROT ID:** Q16891

Full Name:

Mouse Isotype: lgG2b

inner membrane protein, mitochondrial (mitofilin)

Immunogen Catalog Number: AG0102

Calculated MW:

Observed MW:

90 kDa

90 kDa

**Applications** 

**Tested Applications:** 

WB, IF, FC, IHC, Indirect ELISA

Species Specificity:

Human, Rat, Mouse


**Background Information** 

IMMT (also known as mitofilin) is an inner mitochondrial membrane protein that is preferentially expressed in heart tissue and is commonly used as the marker for mitochondria. Mitofilin is known to be a critical organizer of mitochondrial cristae morphology and is indispensable for normal mitochondrial function. Three isoforms of mitofilin exist due to the alternative splicing. All three proteins of 87kD, 89kD and 80kD can be detected in immunoblot analysis with this antibody.


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68226-1-1g (IMMT antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68226-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 68226-1-Ig (IMMT antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 68226-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 68226-1-lg (IMMT antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 68226-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed HeLa cells using IMMT antibody (68226-1-1g, Clone: 2D2E8) at dilution of 1:800 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 68226-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using IMMT antibody (68226-1-Ig, Clone: 2D2E8) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 68226-1-PBS in a different storage buffer formulation.



1X10^6 HEK-293 cells were intracellularly stained with 0.4 ug Anti-Human IMMT (68226-1-Ig, Clone:2D2E8) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2b Isotype Control (66360-3-Ig, Clone: K11B8C4B5) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 68226-1-PBS in a different storage buffer



WB result of IMMT antibody (68226-1-Ig; 1:30000; incubated at room temperature for 1.5 hours) with sh-Control and sh-IMMT transfected HeLa cells. This data was developed using the same antibody clone with 68226-1-PBS in a different storage buffer formulation.