For Research Use Only

HNRNPD Monoclonal antibody

Catalog Number: 68236-1-Ig



Basic Information

Catalog Number: GenBank Accession Number:

68236-1-lg BC026015 Protein A purification

 Size:
 Genel D (NCBI):
 CloneNo.:

 150ul , Concentration: 500 µg/ml by
 3184
 2G9E6

Nanodrop;

Full Name:
Recommended Dilutions:
Source:
heterogeneous nuclear
Mb 1:5000-1:50000
Mouse
ribonucleoprotein D (AU-rich element IHC 1:500-1:2000
RNA binding protein 1, 37kDa)
IF 1:400-1:1600

IgG2bCalculated MW:Immunogen Catalog Number:355 aa, 38 kDaAG3458Observed MW:

38 kDa

Applications

Tested Applications:

IF, IHC, WB, ELISA

Species Specificity:

Human, mouse, rat, pig, rabbit

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A431 cells, LNCaP cells, HeLa cells, HEK-293 cells, Jurkat cells, pig brain tissue, rabbit brain tissue, rat brain tissue, mouse brain tissue

Purification Method:

IHC: human cervical cancer tissue,

IF: HepG2 cells,

Background Information

The heterogeneous nuclear ribonucleoprotein (hnRNP) complexes which provide the substrate for the processing events that pre-mRNAs undergo before becoming functional, translatable mRNAs in the cytoplasm. HNRNPD, also known as AUF1, belongs to the family of AU-binding proteins (AU-BPs) that regulate the cellular half-lives of many mRNAs by directly interacting with an AU-rich element (ARE) located in their 3' untranslated region (PMID:8578590,12704645). AUF1 has four isoforms produced by alternative splicing of a single transcript: p37, p40, p42, and p45 (PMID:9521873).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffe

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68236-1-lg (HNRNPD antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.



Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 68236-1-1g (HNRNPD antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 68236-1-1g (HNRNPD antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using HNRNPD antibody (68236-1-lg, Clone: 2G9E6) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).